Ergod. Th. & Dynam. Sys. (2006), **26**, 1–26
Printed in the United Kingdom © *2006 Cambridge University Press*

arXiv:math/0306136v2 [math.DS] 13 Apr 2006

Asymptotic Randomization of Sofic Shifts by Linear Cellular Automata†

Marcus Pivato and Reem Yassawi
Department of Mathematics, Trent University, 1600 West Bank Drive, Peterborough, Ontario, K9J 7B8, Canada
(e-mail: `marcuspivato@trentu.ca` *and* `ryassawi@trentu.ca`*)*

(*Received July 31, 2004; accepted April 6, 2006*)

Abstract. Let $\mathbb{M} = \mathbb{Z}^D$ be a D-dimensional lattice, and let $(\mathcal{A}, +)$ be an abelian group. $\mathcal{A}^{\mathbb{M}}$ is then a compact abelian group under componentwise addition. A continuous function $\Phi : \mathcal{A}^{\mathbb{M}} \longrightarrow \mathcal{A}^{\mathbb{M}}$ is called a *linear cellular automaton* (LCA) if there is a finite subset $\mathbb{F} \subset \mathbb{M}$ and nonzero coefficients $\varphi_{\mathsf{f}} \in \mathbb{Z}$ so that, for any $\mathbf{a} \in \mathcal{A}^{\mathbb{M}}$, $\Phi(\mathbf{a}) = \sum_{\mathsf{f} \in \mathbb{F}} \varphi_{\mathsf{f}} \cdot \sigma^{\mathsf{f}}(\mathbf{a})$.

Suppose μ is a probability measure on $\mathcal{A}^{\mathbb{M}}$ whose support is a subshift of finite type or sofic shift. We provide sufficient conditions (on Φ and μ) under which Φ *asymptotically randomizes* μ, meaning that $\text{wk}^*\!-\!\lim_{\mathbb{J} \ni j \to \infty} \Phi^j \mu = \eta$, where η is the Haar measure on $\mathcal{A}^{\mathbb{M}}$, and $\mathbb{J} \subset \mathbb{N}$ has Cesàro density 1. In the case when $\Phi = 1 + \sigma$ and $\mathcal{A} = (\mathbb{Z}_{/p})^s$ (p prime), we provide a condition on μ that is both necessary and sufficient. We then use this to construct zero-entropy measures which are randomized by $1 + \sigma$.

MSC: Primary: 37B15; Secondary: 37A50

Let $D \geq 1$, and let $\mathbb{M} := \mathbb{Z}^D$ be the D-dimensional lattice. If $\mathcal{A}$ is a (discretely topologised) finite set, then $\mathcal{A}^{\mathbb{M}}$ is compact in the Tychonoff topology. For any $\mathsf{v} \in \mathbb{M}$, let $\sigma^{\mathsf{v}} : \mathcal{A}^{\mathbb{M}} \longrightarrow \mathcal{A}^{\mathbb{M}}$ be the shift map: $\sigma^{\mathsf{v}}(\mathbf{a}) := \left[b_{\mathsf{m}}|_{\mathsf{m} \in \mathbb{M}} \right]$, where $b_{\mathsf{m}} := a_{\mathsf{m}-\mathsf{v}}$, $\forall \mathsf{m} \in \mathbb{M}$. A *cellular automaton* (CA) is a continuous map $\Phi : \mathcal{A}^{\mathbb{M}} \longrightarrow \mathcal{A}^{\mathbb{M}}$ which commutes with all shifts: for any $\mathsf{m} \in \mathbb{M}$, $\sigma^{\mathsf{m}} \circ \Phi = \Phi \circ \sigma^{\mathsf{m}}$. Let η be the uniform Bernoulli measure on $\mathcal{A}^{\mathbb{M}}$. If μ is another probability measure on $\mathcal{A}^{\mathbb{M}}$, we say Φ *asymptotically randomizes* μ if $\text{wk}^*\!-\!\lim_{\mathbb{J} \ni j \to \infty} \Phi^j \mu = \eta$, where $\mathbb{J} \subset \mathbb{N}$ has Cesàro density one.

If $(\mathcal{A}, +)$ is a finite abelian group, then $\mathcal{A}^{\mathbb{M}}$ is a product group, and η is the Haar measure. A *linear cellular automaton* (LCA) is a CA Φ with a finite subset $\mathbb{F} \subset \mathbb{M}$ (with $\#(\mathbb{F}) \geq 2$), and nonzero coefficients $\varphi_{\mathsf{f}} \in \mathbb{Z}$ (for all $\mathsf{f} \in \mathbb{F}$) so that, for any $\mathbf{a} \in \mathcal{A}^{\mathbb{M}}$,

$$\Phi(\mathbf{a}) \quad = \quad \sum_{\mathsf{f} \in \mathbb{F}} \varphi_{\mathsf{f}} \cdot \sigma^{\mathsf{f}}(\mathbf{a}). \tag{1}$$

---

† This research was partially supported by NSERC Canada, and was also supported by the kind hospitality of the Universidad de Chile during July 2003.

Linear cellular automata are known to asymptotically randomize a wide variety of measures [**MM98, MM99, MHM03, Lin84, FMMN00**], including those satisfying a correlation-decay condition called *harmonic mixing* [**PY02, PY04, MMPY06**]. However, all known sufficient conditions for asymptotic randomization (and for harmonic mixing, in particular) require μ to have *full support*, i.e. $\mathsf{supp}\left(\mu\right) = \mathcal{A}^{\mathbb{M}}$.

We here investigate asymptotic randomization when $\mathsf{supp}\left(\mu\right) \subsetneq \mathcal{A}^{\mathbb{M}}$. In particular we consider the case when $\mathsf{supp}\left(\mu\right)$ is a sofic shift or subshift of finite type. In §1, we let $\mathcal{A} = \mathbb{Z}_{/p}$ (p prime), and demonstrate asymptotic randomization for any Markov random field that is *locally free*, a much weaker assumption than full support. However, in §2 we show that harmonic mixing is a rather restrictive condition, by exhibiting a measure whose support is a mixing sofic shift but which is *not* harmonically mixing.

Thus, in §3, we introduce the less restrictive concept of *dispersion mixing* (for measures) and the dual concept of *dispersion* (for LCA), and state our main result: any dispersive LCA asymptotically randomizes any dispersion mixing measure. In §4, we let $\mathcal{A} = (\mathbb{Z}_{/p})^s$ (p prime, $s \in \mathbb{N}$) and introduce *bipartite* LCA, a broad class exemplified by the automaton $1+\sigma$. We then show that any bipartite LCA is dispersive.

In §5, we show that any *uniformly mixing* and *harmonically bounded* measure is dispersion mixing. In particular, in §6, we show this implies that any mixing Markov measure (supported on a subshift of finite type), and any continuous factor of a mixing Markov measure (supported on a sofic shift) is dispersion mixing, and thus, is asymptotically randomized by any dispersive LCA (e.g. $1+\sigma$). Thus, the example of §2 *is* asymptotically randomized, even though it is not harmonically mixing.

In §7, we refine the results of §3-§4 by introducing *Lucas mixing*, (a weaker condition than dispersion mixing). When $\mathcal{A} = (\mathbb{Z}_{/p})^s$, we show that a measure is asymptotically randomized by the automaton $1+\sigma$ if and *only if* it is Lucas mixing. Finally, in §8, we use Lucas mixing to construct a class of *zero-entropy* measures which are asymptotically randomized by randomized by $1+\sigma$, thereby refuting the conjecture that positive entropy is necessary for asymptotic randomization.

Preliminaries & Notation: Throughout, $(\mathcal{A}, +)$ is an abelian group (usually $\mathcal{A} = (\mathbb{Z}_{/p})^s$,where p is prime and $s \in \mathbb{N}$). Elements of $\mathcal{A}^{\mathbb{M}}$ are denoted by boldfaced letters (e.g. $\mathbf{a}$, $\mathbf{b}$, $\mathbf{c}$), and subsets by gothic letters (e.g. $\mathfrak{A}$, $\mathfrak{B}$, $\mathfrak{C}$). Elements of $\mathbb{M}$ are sans serif (e.g. l, m, n) and subsets are $\mathbb{U}, \mathbb{V}, \mathbb{W}$.

If $\mathbb{U} \subset \mathbb{M}$ and $\mathbf{a} \in \mathcal{A}^{\mathbb{M}}$ then $\mathbf{a}_{\mathbb{U}} := \left[a_{\mathsf{u}}|_{\mathsf{u}\in\mathbb{U}}\right]$ is the 'restriction' of $\mathbf{a}$ to an element of $\mathcal{A}^{\mathbb{U}}$. For any $\mathbf{b} \in \mathcal{A}^{\mathbb{U}}$, let $[\mathbf{b}] := \left\{\mathbf{c} \in \mathcal{A}^{\mathbb{M}} \,;\, \mathbf{c}_{\mathbb{U}} = \mathbf{b}\right\}$ be the corresponding cylinder set. In particular, if $\mathbf{a} \in \mathcal{A}^{\mathbb{M}}$, then $[\mathbf{a}_{\mathbb{U}}] := \left\{\mathbf{c} \in \mathcal{A}^{\mathbb{M}} \,;\, \mathbf{c}_{\mathbb{U}} = \mathbf{a}_{\mathbb{U}}\right\}$.

Measures: Let $\mathcal{M}(\mathcal{A}^{\mathbb{M}})$ be the set of Borel probability measures on $\mathcal{A}^{\mathbb{M}}$. If $\mu \in \mathcal{M}(\mathcal{A}^{\mathbb{M}})$ and $\mathbb{I} \subset \mathbb{M}$, then let $\mu_{\mathbb{I}} \in \mathcal{M}(\mathcal{A}^{\mathbb{I}})$ be the marginal projection of μ onto $\mathcal{A}^{\mathbb{I}}$. If $\mathbb{J} \subset \mathbb{M}$ and $\mathbf{b} \in \mathcal{A}^{\mathbb{J}}$, then let $\mu^{(\mathbf{b})} \in \mathcal{M}(\mathcal{A}^{\mathbb{M}})$ be the conditional probability measure in the cylinder set $[\mathbf{b}]$. In other words, for any $\mathfrak{X} \subset \mathcal{A}^{\mathbb{M}}$, $\mu^{(\mathbf{b})}[\mathfrak{X}] := \mu\left(\mathfrak{X} \cap [\mathbf{b}]\right)/\mu\,[\mathbf{b}]$. In particular, if $\mathbb{I} \subset \mathbb{M}$ is finite, then $\mu^{(\mathbf{b})}_{\mathbb{I}} \in \mathcal{M}(\mathcal{A}^{\mathbb{M}})$ is the conditional probability measure on the $\mathbb{I}$ coordinates: for any $\mathbf{c} \in \mathcal{A}^{\mathbb{I}}$, $\mu^{(\mathbf{b})}_{\mathbb{I}}[\mathbf{c}] := \mu\left([\mathbf{c}] \cap [\mathbf{b}]\right)/\mu\,[\mathbf{b}]$.

Subshifts: A *subshift* [**Kit98, LM95**] is a a closed, shift-invariant subset $\mathfrak{X} \subset \mathcal{A}^{\mathbb{M}}$. If $\mathbb{U} \subset \mathbb{M}$, then let $\mathfrak{X}_{\mathbb{U}} := \{\mathbf{x}_{\mathbb{U}} \ ; \ \mathbf{x} \in \mathfrak{X}\}$ be all *admissible* $\mathbb{U}$-*blocks* in $\mathfrak{X}$. If $\mathbb{U} \subset \mathbb{M}$ is finite, and $\mathfrak{W} = \{\mathbf{w}_1, \ldots, \mathbf{w}_N\} \subset \mathcal{A}^{\mathbb{U}}$ is a collection of admissible blocks, then the induced *subshift of finite type* (SFT) is the largest subshift $\mathfrak{X} \subset \mathcal{A}^{\mathbb{M}}$ such that $\mathfrak{X}_{\mathbb{U}} = \mathfrak{W}$. In other words, $\mathfrak{X} := \bigcap_{\mathsf{m} \in \mathbb{M}} \sigma^{\mathsf{m}} [\mathfrak{W}]$, where $[\mathfrak{W}] := \left\{\mathbf{a} \in \mathcal{A}^{\mathbb{M}} \ ; \ \mathbf{a}_{\mathbb{U}} \in \mathfrak{W}\right\}$. A *sofic shift* is the image of an SFT under a block map.

In particular, if $\mathbb{M} = \mathbb{Z}$ and $\mathbb{U} = \{0, 1\}$, then $\mathfrak{X}$ is called *topological Markov shift*, and the *transition matrix* of $\mathfrak{X}$ is the matrix $\mathbf{P} = [p_{ab}]_{a,b \in \mathcal{A}}$, where $p_{ab} = 1$ if $[ab] \in \mathfrak{W}$, and $p_{ab} = 0$ if $[ab] \notin \mathfrak{W}$.

Characters: Let $\mathbb{T}^1 \subset \mathbb{C}$ be the circle group. A *character* of $\mathcal{A}^{\mathbb{M}}$ is a continuous homomorphism $\boldsymbol{\chi} : \mathcal{A}^{\mathbb{M}} \longrightarrow \mathbb{T}^1$; the group of such characters is denoted $\widehat{\mathcal{A}^{\mathbb{M}}}$. For any $\boldsymbol{\chi} \in \widehat{\mathcal{A}^{\mathbb{M}}}$ there is a finite subset $\mathbb{K} \subset \mathbb{M}$, and nontrivial $\chi_{\mathsf{k}} \in \widehat{\mathcal{A}}$ for all $\mathsf{k} \in \mathbb{K}$, such that, for any $\mathbf{a} \in \mathcal{A}^{\mathbb{M}}$, $\boldsymbol{\chi}(\mathbf{a}) = \prod_{\mathsf{k} \in \mathbb{K}} \chi_{\mathsf{k}}(a_{\mathsf{k}})$. We indicate this by writing: "$\boldsymbol{\chi} = \bigotimes_{\mathsf{k} \in \mathbb{K}} \chi_{\mathsf{k}}$". The *rank* of $\boldsymbol{\chi}$ is the cardinality of $\mathbb{K}$.

Cesàro Density: If $\ell, n \in \mathbb{Z}$, then let $[\ell...n) := \{m \in \mathbb{Z} \ ; \ \ell \leq m < n\}$. If $\mathbb{J} \subset \mathbb{N}$, then the *Cesàro density* of $\mathbb{J}$ is defined: $\mathsf{density}\,(\mathbb{J}) := \lim_{N \to \infty} \frac{1}{N} \# \Big(\mathbb{J} \cap [0...N)\Big)$. If $\mathbb{J}, \mathbb{K} \subset \mathbb{N}$, then their *relative Cesàro density* is defined:

$$\mathsf{rel\,density}\,[\mathbb{J}/\mathbb{K}] \quad := \quad \lim_{N \to \infty} \frac{\#\left(\mathbb{J} \cap [0...N)\right)}{\#\left(\mathbb{K} \cap [0...N)\right)}.$$

In particular, $\mathsf{density}\,(\mathbb{J}) = \mathsf{rel\,density}\,[\mathbb{J}/\mathbb{N}]$.

1. *Harmonic Mixing of Markov Random Fields*

Let $\mathbb{B} \subset \mathbb{M}$ be a finite subset, symmetric under multiplication by -1 (usually, $\mathbb{B} = \{-1, 0, 1\}^D$). For any $\mathbb{U} \subset \mathbb{M}$, we define

$$\mathsf{cl}\,(\mathbb{U}) \ := \ \{\mathsf{u} + \mathsf{b} \ ; \ \mathsf{u} \in \mathbb{U} \text{ and } \ \mathsf{b} \in \mathbb{B}\} \qquad \text{and} \qquad \partial\mathbb{U} \ := \ \mathsf{cl}\,(\mathbb{U}) \setminus \mathbb{U}.$$

For example, if $\mathbb{M} = \mathbb{Z}$ and $\mathbb{B} = \{-1, 0, 1\}$, then $\partial\{0\} = \{\pm 1\}$.

Let $\mu \in \mathcal{M}(\mathcal{A}^{\mathbb{M}})$. Suppose $\mathbb{U} \subset \mathbb{M}$, and let $\mathbb{V} := \partial\mathbb{U}$ and $\mathbb{W} = \mathbb{M} \setminus \mathsf{cl}\,(\mathbb{U})$. If $\mathbf{b} \in \mathcal{A}^{\mathbb{V}}$, then we say $\mathbf{b}$ *isolates* $\mathbb{U}$ from $\mathbb{W}$ if the conditional measure $\mu^{(\mathbf{b})}$ is a product of $\mu^{(\mathbf{b})}_{\mathbb{U}}$ and $\mu^{(\mathbf{b})}_{\mathbb{W}}$. That is, for any $\mathfrak{U} \subset \mathcal{A}^{\mathbb{U}}$ and $\mathfrak{W} \subset \mathcal{A}^{\mathbb{W}}$, we have $\mu^{(\mathbf{b})}\left(\mathfrak{U} \cap \mathfrak{W}\right) = \mu^{(\mathbf{b})}_{\mathbb{U}}(\mathfrak{U}) \cdot \mu^{(\mathbf{b})}_{\mathbb{W}}(\mathfrak{W})$.

We say that μ is a *Markov random field* [**Bré99, KS80**] with *interaction range* $\mathbb{B}$ (or write, "μ is a $\mathbb{B}$-*MRF*") if, for any $\mathbb{U} \subset \mathbb{M}$ with $\mathbb{V} = \partial\mathbb{U}$ and $\mathbb{W} = \mathbb{M} \setminus \mathsf{cl}\,(\mathbb{U})$, any choice of $\mathbf{b} \in \mathcal{A}^{\mathbb{V}}$ isolates $\mathbb{U}$ from $\mathbb{W}$.

For example, if $\mathbb{M} = \mathbb{Z}$ and $\mathbb{B} = \{-1, 0, 1\}$, then μ is a $\mathbb{B}$-MRF iff μ is a (one-step) Markov chain. If $\mathbb{B} = [-N...N]$, then μ is a $\mathbb{B}$-MRF iff μ is an N-step Markov chain.

LEMMA 1.1. *If μ is a Markov random field, then $\mathsf{supp}\,(\mu)$ is a subshift of finite type.* □

For example, if μ is a Markov chain on $\mathcal{A}^{\mathbb{Z}}$, then $\mathsf{supp}\,(\mu)$ is a topological Markov shift.

Let $\mathbb{B} \subset \mathbb{M}$, and let $\mu \in \mathcal{M}(\mathcal{A}^{\mathbb{M}})$ be $\mathbb{B}$-MRF. Let $\mathbb{S} := \mathbb{B} \setminus \{0\}$. For any $\mathbf{b} \in \mathcal{A}^{\mathbb{S}}$, let $\mu^{(\mathbf{b})}_0 \in \mathcal{M}(\mathcal{A})$ be the conditional probability measure on the zeroth coordinate. We say that μ is *locally free* if, for any $\mathbf{b} \in \mathcal{A}^{\mathbb{S}}$, $\#\left(\mathsf{supp}\left(\mu^{(\mathbf{b})}_0\right)\right) \geq 2$.

EXAMPLE: If $D = 1$, then $\mathbb{B} = \{-1, 0, 1\}$, $\mathbb{S} = \{\pm 1\}$, and μ is a Markov chain. Thus, $\mathsf{supp}(\mu)$ is a topological Markov shift, with transition matrix $\mathbf{P} = [p_{ab}]_{a,b\in\mathcal{A}}$. For any $a, b \in \mathcal{A}$, write $a \rightsquigarrow b$ if $p_{ab} = 1$, and define the *follower* and *predecessor* sets

$$\mathcal{F}(a) \ := \ \{b \in \mathcal{A} \,;\, a \rightsquigarrow b\} \quad \text{and} \quad \mathcal{P}(b) \ := \ \{a \in \mathcal{A} \,;\, a \rightsquigarrow b\}.$$

It is easy to show that the following are equivalent:

1. μ is locally free.
2. Every entry of $\mathbf{P}^2$ is 2 or larger.
3. For any $a, b \in \mathcal{A}$, $\#(\mathcal{F}(a) \cap \mathcal{P}(b)) \geq 2$.

Recall that $\widehat{\mathcal{A}}$ is the dual group of $\mathcal{A}$. For any $\boldsymbol{\chi} \in \widehat{\mathcal{A}}$ and $\nu \in \mathcal{M}(\mathcal{A})$, let $\langle \chi, \nu \rangle := \sum_{a\in\mathcal{A}} \chi(a) \cdot \nu\{a\}$. It is easy to check:

LEMMA 1.2. *Let p be prime and $\mathcal{A} = \mathbb{Z}_{/p}$. If μ is a locally free MRF on $\mathcal{A}^{\mathbb{M}}$, then there is some $c < 1$ such that, for all nontrivial $\chi \in \widehat{\mathcal{A}}$, and any $\mathbf{b} \in \mathcal{A}^{\mathbb{S}}$,* $\left|\left\langle \chi, \mu_0^{(\mathbf{b})} \right\rangle\right| \ \leq \ c.$ □

For any $\boldsymbol{\chi} \in \widehat{\mathcal{A}^{\mathbb{M}}}$ and $\mu \in \mathcal{M}\left(\mathcal{A}^{\mathbb{M}}\right)$, define $\langle \boldsymbol{\chi}, \mu \rangle \ := \ \int_{\mathcal{A}^{\mathbb{M}}} \boldsymbol{\chi}(\mathbf{a}) \ d\mu[\mathbf{a}]$. A measure μ is called *harmonically mixing* if, for any $\epsilon > 0$, there is some $R \in \mathbb{N}$ such that, for any $\boldsymbol{\chi} \in \widehat{\mathcal{A}^{\mathbb{M}}}$,

$$\Big(\mathsf{rank}\,[\boldsymbol{\chi}] \ > \ R\Big) \Longrightarrow \Big(|\langle \boldsymbol{\chi}, \mu \rangle| \ < \ \epsilon\Big).$$

The significance of this is the following [**PY02**, Theorem 12]:

THEOREM: *Let $\mathcal{A} = \mathbb{Z}_{/p}$, where p is prime. Any LCA on $\mathcal{A}^{\mathbb{M}}$ asymptotically randomizes any harmonically mixing measure.* □

Most MRFs with full support are harmonically mixing [**PY04**, Theorem 15]. We now extend this.

THEOREM 1.3. *Let $\mathcal{A} = \mathbb{Z}_{/p}$, where p is prime. Any locally free MRF on $\mathcal{A}^{\mathbb{M}}$ is harmonically mixing.*

Proof. Let μ be a locally free $\mathbb{B}$-MRF. A subset $\mathbb{I} \subset \mathbb{M}$ is $\mathbb{B}$-*separated* if $(\mathsf{i} - \mathsf{j}) \not\in \mathbb{B}$ for all $\mathsf{i}, \mathsf{j} \in \mathbb{I}$ with $\mathsf{i} \neq \mathsf{j}$. Let $\mathbb{K} \subset \mathbb{M}$ be finite, and let $\boldsymbol{\chi} \ := \ \bigotimes_{\mathsf{k}\in\mathbb{K}} \chi_{\mathsf{k}}$ be a character of $\mathcal{A}^{\mathbb{M}}$.

CLAIM 1: *Let $K := \#(\mathbb{K}) = \mathsf{rank}\,[\boldsymbol{\chi}]$, and let $B \ := \ \max\{|\mathsf{b}_1 - \mathsf{b}_2| \,;\, \mathsf{b}_1, \mathsf{b}_2 \in \mathbb{B}\}$. There exists a $\mathbb{B}$-separated subset $\mathbb{I} \subset \mathbb{K}$ such that*

$$\#(\mathbb{I}) \quad = \quad I \quad \geq \quad \frac{K}{B^D}. \tag{2}$$

Proof. Let $\widetilde{\mathbb{B}} := [0...B)^D$ be a box of sidelength B. Cover $\mathbb{K}$ with disjoint translated copies of $\widetilde{\mathbb{B}}$, so that

$$\mathbb{K} \quad \subset \quad \bigsqcup_{\mathsf{i}\in\mathbb{I}} \left(\widetilde{\mathbb{B}} + \mathsf{i}\right)$$

for some set $\mathbb{I} \subset \mathbb{K}$. Thus, $|\mathsf{i} - \mathsf{j}| \geq B$ for any $\mathsf{i}, \mathsf{j} \in \mathbb{I}$ with $\mathsf{i} \neq \mathsf{j}$, so $(\mathsf{i} - \mathsf{j}) \not\in \mathbb{B}$. Also, $\#\left(\widetilde{\mathbb{B}}\right) = B^D$, so each copy covers at most B^D points in $\mathbb{K}$. Thus, we require at least $\frac{K}{B^D}$ copies to cover all of $\mathbb{K}$. In other words, $I \geq \frac{K}{B^D}$. ◇ `Claim 1`

Thus, $\boldsymbol{\chi} = \boldsymbol{\chi}_{\mathbb{I}} \cdot \boldsymbol{\chi}_{\mathbb{K}\setminus\mathbb{I}}$, where $\boldsymbol{\chi}_{\mathbb{I}}(\mathbf{a}) := \prod_{\mathsf{i}\in\mathbb{I}} \chi_{\mathsf{i}}(a_{\mathsf{i}})$, and $\boldsymbol{\chi}_{\mathbb{K}\setminus\mathbb{I}}(\mathbf{a}) := \prod_{\mathsf{k}\in\mathbb{K}\setminus\mathbb{I}} \chi_{\mathsf{k}}(a_{\mathsf{k}})$.

Let $\mathbb{J} := (\partial\,\mathbb{I}) \cup (\mathbb{K}\setminus\mathbb{I})$; fix $\mathbf{b} \in \mathcal{A}^{\mathbb{J}}$, and let $\mu_{\mathbb{I}}^{(\mathbf{b})} \in \mathcal{M}(\mathcal{A}^{\mathbb{I}})$ be the corresponding conditional probability measure. Since μ is a Markov random field, and the $\mathbb{I}$ coordinates are 'isolated' from one another by $\mathbb{J}$ coordinates, it follows that $\mu_{\mathbb{I}}^{(\mathbf{b})}$ is a product measure. In other words, for any $\mathbf{a} \in \mathcal{A}^{\mathbb{I}}$,

$$\mu_{\mathbb{I}}^{(\mathbf{b})}[\mathbf{a}] \quad = \quad \prod_{\mathsf{i}\in\mathbb{I}} \mu_{\mathsf{i}}^{(\mathbf{b})}\{a_{\mathsf{i}}\}. \tag{3}$$

Thus, the conditional expectation of $\boldsymbol{\chi}_{\mathbb{I}}$ is given:

$$\begin{aligned}\left\langle \boldsymbol{\chi}_{\mathbb{I}},\ \mu_{\mathbb{I}}^{(\mathbf{b})} \right\rangle \quad &= \quad \sum_{\mathbf{a}\in\mathcal{A}^{\mathbb{I}}} \mu_{\mathbb{I}}^{(\mathbf{b})}[\mathbf{a}] \cdot \left(\prod_{\mathsf{i}\in\mathbb{I}} \chi_{\mathsf{i}}(a_{\mathsf{i}}) \right) \quad \underset{(*)}{=\!=} \quad \sum_{\mathbf{a}\in\mathcal{A}^{\mathbb{I}}} \left(\prod_{\mathsf{i}\in\mathbb{I}} \mu_{\mathsf{i}}^{(\mathbf{b})}\{a_{\mathsf{i}}\} \cdot \chi_{\mathsf{i}}(a_{\mathsf{i}}) \right) \\ &= \quad \prod_{\mathsf{i}\in\mathbb{I}} \left(\sum_{a_{\mathsf{i}}\in\mathcal{A}} \mu^{(\mathbf{b})}\{a_{\mathsf{i}}\} \cdot \chi_{\mathsf{i}}(a_{\mathsf{i}}) \right) \quad = \quad \prod_{\mathsf{i}\in\mathbb{I}} \left\langle \chi_{\mathsf{i}},\ \mu_{\mathsf{i}}^{(\mathbf{b})} \right\rangle,\end{aligned}$$

where $(*)$ is by equation (3). Thus, $\left\langle \boldsymbol{\chi},\ \mu^{(\mathbf{b})} \right\rangle = \boldsymbol{\chi}_{\mathbb{K}\setminus\mathbb{I}}(\mathbf{b}) \cdot \left\langle \boldsymbol{\chi}_{\mathbb{I}},\ \mu_{\mathbb{I}}^{(\mathbf{b})} \right\rangle = \boldsymbol{\chi}_{\mathbb{K}\setminus\mathbb{I}}(\mathbf{b}) \cdot \prod_{\mathsf{i}\in\mathbb{I}} \left\langle \chi_{\mathsf{i}},\ \mu_{\mathsf{i}}^{(\mathbf{b})} \right\rangle$. Thus, if $I = \#(\mathbb{I})$, then

$$\left| \left\langle \boldsymbol{\chi},\ \mu^{(\mathbf{b})} \right\rangle \right| \quad = \quad \left| \boldsymbol{\chi}_{\mathbb{K}\setminus\mathbb{I}}(\mathbf{b}) \right| \cdot \prod_{\mathsf{i}\in\mathbb{I}} \left| \left\langle \chi_{\mathsf{i}},\ \mu_{\mathsf{i}}^{(\mathbf{b})} \right\rangle \right| \quad \leq \quad 1 \cdot c^{I} \tag{4}$$

where the last step follows from Lemma 1.2. But $\langle \boldsymbol{\chi}, \mu \rangle = \sum_{\mathbf{b}\in\mathcal{A}^{\mathbb{J}}} \mu[\mathbf{b}] \cdot \left\langle \boldsymbol{\chi},\ \mu^{(\mathbf{b})} \right\rangle$, so

$$|\langle \boldsymbol{\chi}, \mu \rangle| \quad \leq \quad \sum_{\mathbf{b}\in\mathcal{A}^{\mathbb{J}}} \mu[\mathbf{b}] \cdot \left| \left\langle \boldsymbol{\chi},\ \mu^{(\mathbf{b})} \right\rangle \right| \quad \underset{(*)}{\leq} \quad \sum_{\mathbf{b}\in\mathcal{A}^{\mathbb{J}}} \mu[\mathbf{b}] \cdot c^{I} \quad = \quad c^{I} \quad \underset{(\dagger)}{\leq} \quad c^{K/(B^D)} \xrightarrow[K\to\infty]{} 0.$$

Here $(*)$ is by equation (4) and $(\dagger)$ is by equation (2). □

2. *The Even Shift is Not Harmonically Mixing*

We will now construct a measure ν, supported on a sofic shift, which is *not* harmonically mixing. Nonetheless, we'll show in §3-§5 that this measure *is* asymptotically randomized by many LCA.

Let $\mathfrak{X} \subset \left(\mathbb{Z}_{/3}\right)^{\mathbb{Z}}$ be the subshift of finite type defined by the transition matrix

$$\mathbf{A} \quad = \quad \begin{bmatrix} 1 & 0 & 1 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}, \quad \text{where, } \forall i, j \in \mathbb{Z}_{/3}, \quad a_{ij} = \begin{cases} 1 & \text{if } j \rightsquigarrow i \text{ is allowed} \\ 0 & \text{if } j \rightsquigarrow i \text{ is not allowed} \end{cases}$$

Let $\Phi : \mathfrak{X} \to \left(\mathbb{Z}_{/2}\right)^{\mathbb{Z}}$ be the factor map of radius 0 which sends 0 into 0 and both 1 and 2 to 1. Then $\mathfrak{S} := \Phi(\mathfrak{X})$ is Weiss's *Even Sofic Shift*: if $\mathbf{s} \in \mathfrak{S}$, then there are an even number of 1's between any two occurrences of 0 in $\mathbf{s}$.

For any $N \in \mathbb{N}$, and $i, j \in \mathbb{Z}_{/3}$, let $\mathfrak{X}_{ij}^{N} := \{\mathbf{x} \in \mathfrak{X}\ ;\ x_0 = i,\ x_N = j\}$, and let:

$$\mathfrak{E}_N \ := \ \left\{ \mathbf{s} \in \mathfrak{S}\ ;\ \sum_{n=0}^{N} s_n \text{ is even} \right\}, \ \text{and } \mathfrak{O}_N \ := \ \left\{ \mathbf{s} \in \mathfrak{S}\ ;\ \sum_{n=0}^{N} s_n \text{ is odd} \right\}.$$

Lemma 2.1. *$\forall i, j \in \mathbb{Z}_{/3}$, either $\Phi\left(\mathfrak{X}^N_{i,j}\right) \subset \mathfrak{E}_N$ or $\Phi\left(\mathfrak{X}^N_{i,j}\right) \subset \mathfrak{O}_N$. In particular,*

$$\Phi\left(\mathfrak{X}^N_{0,0} \sqcup \mathfrak{X}^N_{1,2} \sqcup \mathfrak{X}^N_{2,1} \sqcup \mathfrak{X}^N_{0,2} \sqcup \mathfrak{X}^N_{1,0}\right) \;=\; \mathfrak{E}_N \qquad \textit{and} \qquad \Phi\left(\mathfrak{X}^N_{1,1} \sqcup \mathfrak{X}^N_{0,1} \sqcup \mathfrak{X}^N_{2,0} \sqcup \mathfrak{X}^N_{2,2}\right) \;=\; \mathfrak{O}_N.$$

Proof. Let $\mathbf{x} \in \mathfrak{X}^N_{ij}$, and $\mathbf{s} := \Phi(\mathbf{x})$. Note that, if $k < k^*$ are any two values such that $x_k = 0 = x_{k^*}$, then $\sum_{n=k}^{k^*} s_n$ is even. In particular, let k be the first element of $[0...N]$ where $x_k = 0$, and let k^* be the last element of $[0...N]$ where $x_{k^*} = 0$. Thus, $\sum_{n=k}^{k^*} s_n \equiv 0 \pmod 2$, so that $\sum_{n=0}^{N} s_n \;\equiv\; \sum_{n=0}^{k-1} s_n + \sum_{n=k^*+1}^{N} s_n \pmod 2$.

But since $x_{k-1} \neq 0 \neq x_{k^*+1}$ by construction, the definition of $\mathfrak{X}$ forces $x_{k-1} = 2$ and $x_{k^*+1} = 1$. Thus the parity of $\sum_{n=0}^{k-1} s_n$ depends only on the value of $x_0 = i$. Similarly the parity of $\sum_{n=k^*+1}^{N} s_n$ depends only on $x_N = j$. □

Let $\mu \in \mathcal{M}[\mathfrak{X}]$ be a mixing Markov measure on $\mathfrak{X}$, with transition matrix $\mathbf{P}$ and Perron measure $\boldsymbol{\rho} = (\rho_0, \rho_1, \rho_2) \in \mathcal{M}\left[\mathbb{Z}_{/3}\right]$. Let $\nu := \Phi\mu \in \mathcal{M}[\mathfrak{S}]$, so that if $\mathfrak{U} \subset \mathfrak{S}$ is measurable, then $\nu[\mathfrak{U}] := \mu\left[\Phi^{-1}(\mathfrak{U})\right]$.

For all $N \in \mathbb{N}$, define character $\boldsymbol{\chi}_N$ by $\boldsymbol{\chi}_N(\mathbf{x}) := \prod_{n=0}^{N} (-1)^{x_n}$ for all $\mathbf{x} \in \left(\mathbb{Z}_{/2}\right)^{\mathbb{Z}}$. Then Lemma 2.1 implies:

$$\begin{aligned} \langle \boldsymbol{\chi}_N, \nu \rangle &= \nu(\mathfrak{E}_N) - \nu(\mathfrak{O}_N) \\ &= \mu\left(\mathfrak{X}^N_{0,0} \sqcup \mathfrak{X}^N_{1,2} \sqcup \mathfrak{X}^N_{2,1} \sqcup \mathfrak{X}^N_{0,2} \sqcup \mathfrak{X}^N_{1,0}\right) \;-\; \nu\left(\mathfrak{X}^N_{1,1} \sqcup \mathfrak{X}^N_{0,1} \sqcup \mathfrak{X}^N_{2,0} \sqcup \mathfrak{X}^N_{2,2}\right). \end{aligned}$$

But μ is mixing, so $\lim_{N\to\infty} \mu(\mathfrak{X}^N_{i,j}) \;=\; \rho_i \cdot \rho_j$. Thus, $\lim_{N\to\infty} \langle \boldsymbol{\chi}_N, \nu \rangle \;=\; \rho_0^2 + 2\rho_1\rho_2 - \rho_1^2 - \rho_2^2$. So for example if

$$\mathbf{P} = \begin{bmatrix} 1/2 & 0 & 1/2 \\ 1/2 & 0 & 1/2 \\ 0 & 1 & 0 \end{bmatrix}$$

with Perron measure $\boldsymbol{\rho} \;=\; \left(\frac{2}{5}, \frac{1}{5}, \frac{2}{5}\right)$, then $\lim_{N\to\infty} \langle \boldsymbol{\chi}_N, \nu \rangle \neq 0$. But clearly, $\mathsf{rank}\left[\boldsymbol{\chi}_N\right] = N$, so that $\lim_{N\to\infty} \mathsf{rank}\left[\boldsymbol{\chi}_N\right] \;=\; \infty$. Thus ν is not harmonically mixing.

3. *Dispersion Mixing*

The example from §2 suggests the need for an asymptotic randomization condition on measures that is less restrictive than harmonic mixing. In this section, we'll define the concepts of *dispersion mixing* (for measures) and *dispersion* (for automata) which together yield asymptotic randomization. In §4 we'll show that many LCA are dispersive. In §5 and §6 we'll show that many measures (including the Even Shift measure ν from §2) are dispersion mixing.

Let Φ be an LCA as in equation (1). The advantage of this 'polynomial' notation is that composition of two LCA corresponds to multiplication of their respective polynomials. For example, suppose $\mathcal{A} = (\mathbb{Z}_{/p})^s$, where $p \in \mathbb{N}$ is prime, and $s \in \mathbb{N}$. Suppose $\mathbb{M} = \mathbb{Z}$ and $\Phi = 1 + \sigma$; that is, $\Phi(\mathbf{a})_0 = a_0 + a_1 \pmod p$. Then the Binomial Theorem implies:

$$\text{For any } N \in \mathbb{N}, \qquad \Phi^N \quad = \quad \sum_{n=0}^{N} \begin{bmatrix} N \\ n \end{bmatrix}_p \sigma^n, \qquad \text{where} \quad \begin{bmatrix} N \\ n \end{bmatrix}_p \;:=\; \binom{N}{n} \bmod p. \tag{5}$$

Let $S > 0$, and let $\mathbb{K}, \mathbb{J} \subset \mathbb{M}$ be subsets. We say that $\mathbb{K}$ and $\mathbb{J}$ are S-*separated* if

$$\min\{|\mathsf{k} - \mathsf{j}| \;;\; \mathsf{k} \in \mathbb{K} \text{ and } \mathsf{j} \in \mathbb{J}\} \quad \geq \quad S$$

If $\mathbb{F}, \mathbb{G} \subset \mathbb{M}$, and $\Phi \;=\; \sum_{\mathsf{f}\in\mathbb{F}} \varphi_\mathsf{f} \cdot \sigma^\mathsf{f}$ and $\Gamma \;=\; \sum_{\mathsf{g}\in\mathbb{G}} \gamma_\mathsf{g} \cdot \sigma^\mathsf{g}$ are two LCA, then we say Φ and Γ are S-*separated* if $\mathbb{F}$ and $\mathbb{G}$ are S-separated. Likewise, if $\mathbb{K}, \mathbb{X} \subset \mathbb{M}$, and $\boldsymbol{\chi} \;=\; \bigotimes_{\mathsf{k}\in\mathbb{K}} \chi_\mathsf{k}$ and $\boldsymbol{\xi} \;=\; \bigotimes_{\mathsf{x}\in\mathbb{X}} \xi_\mathsf{x}$ are two characters, then we say $\boldsymbol{\chi}$ and $\boldsymbol{\xi}$ are S-*separated* if $\mathbb{K}$ and $\mathbb{X}$ are S-separated.

If $\Phi \;=\; \sum_{\mathsf{f}\in\mathbb{F}} \varphi_\mathsf{f} \cdot \sigma^\mathsf{f}$ is an LCA, then let $\mathsf{rank}_S(\Phi)$ be the maximum number of S-separated LCA which can be summed to yield Φ. That is:

$$\mathsf{rank}_S(\Phi) \quad := \quad \max\Big\{R \;;\; \exists\, \Phi_1, \ldots, \Phi_R \text{ mutually } S\text{-separated, with } \Phi \;=\; \Phi_1 + \cdots + \Phi_R\Big\}.$$

For example, if

$$\Phi \quad = \quad 1 \;+\; \sigma^5 + \sigma^6 \;+\; \sigma^{11} + \sigma^{12} + \sigma^{13},$$

then $\mathsf{rank}_4(\Phi) = 3$, because $\Phi = \Phi_1 + \Phi_2 + \Phi_3$, where

$$\Phi_1 \;=\; 1, \quad \Phi_2 \;=\; \sigma^5 + \sigma^6, \text{ and } \Phi_3 \;=\; \sigma^{11} + \sigma^{12} + \sigma^{13}.$$

On the other hand, clearly, $\mathsf{rank}_1(\Phi) = 6$, while $\mathsf{rank}_7(\Phi) = 1$.

Likewise, if $\boldsymbol{\chi} \;=\; \bigotimes_{\mathsf{k}\in\mathbb{K}} \chi_\mathsf{k}$ is a character, and $S > 0$, then we define

$$\mathsf{rank}_S(\chi) \quad := \quad \max\Big\{R \;;\; \exists\, \boldsymbol{\chi}_1, \ldots, \boldsymbol{\chi}_R \text{ mutually } S\text{-separated, with } \boldsymbol{\chi} = \boldsymbol{\chi}_1 \otimes \cdots \otimes \boldsymbol{\chi}_R\Big\}.$$

(In the notation of §1, $\mathsf{rank}[\boldsymbol{\chi}] \;=\; \mathsf{rank}_1(\boldsymbol{\chi})$.)

We say that μ is *dispersion mixing* (DM) if, for every $\epsilon > 0$, there exist $S, R > 0$ such that, for any character $\boldsymbol{\chi} \in \widehat{\mathcal{A}^\mathbb{M}}$, $\Big(\mathsf{rank}_S(\boldsymbol{\chi}) \;>\; R\Big) \Longrightarrow \Big(|\langle\boldsymbol{\chi}, \mu\rangle| \;<\; \epsilon\Big)$. Note that dispersion mixing is less restrictive than harmonic mixing.

If Φ is an LCA and $\boldsymbol{\chi}$ is a character, then $\boldsymbol{\chi} \circ \Phi$ is also a character. We say that Φ is *dispersive* if, for any $S > 0$, and any character $\boldsymbol{\chi} \in \widehat{\mathcal{A}^\mathbb{M}}$, there is a subset $\mathbb{J} \subset \mathbb{N}$ of density 1 such that $\lim_{\mathbb{J}\ni j\to\infty} \mathsf{rank}_S\left(\boldsymbol{\chi} \circ \Phi^j\right) \;=\; \infty$. It follows:

THEOREM 3.1. *Let $\mathcal{A}$ be any finite abelian group. If $\Phi : \mathcal{A}^\mathbb{M} \longrightarrow \mathcal{A}^\mathbb{M}$ is a dispersive LCA and $\mu \in \mathcal{M}(\mathcal{A}^\mathbb{M})$ is dispersion mixing, then Φ asymptotically randomizes μ.* □

Theorem 3.1 is an immediate consequence of an easily verified lemma:

Lemma 3.2. *Φ asymptotically randomizes μ if and only if, for all $\boldsymbol{\chi} \in \widehat{\mathcal{A}^{\mathbb{M}}}$, there is a subset $\mathbb{J} \subset \mathbb{N}$ with $\mathsf{density}(\mathbb{J}) = 1$, such that* $\displaystyle\lim_{\mathbb{J}\ni j\to\infty} \left|\langle \boldsymbol{\chi}\circ\Phi^j,\ \mu\rangle\right| \quad = \quad 0.$

Proof. See the proof of Theorem 12 in [**PY02**]. □

4. *Dispersion and Bipartite CA*

If $\mathsf{m} = (m_1, m_2, \ldots, m_D) \in \mathbb{M}$, then let $|\mathsf{m}| := |m_1| + |m_2| + \cdots + |m_D|$. If $\Gamma \;=\; \sum_{\mathsf{g}\in\mathbb{G}} \gamma_{\mathsf{g}}\cdot\sigma^{\mathsf{g}}$ is a linear cellular automaton, then define $\mathsf{diam}[\Gamma] \;:=\; \max\{|\mathsf{g}-\mathsf{h}|\ ;\ \mathsf{g},\mathsf{h}\in\mathbb{G}\}$.

The *centre* of Γ is the centroid of $\mathbb{G}$ (as a subset of $\mathbb{R}^n$):

$$\mathsf{centre}(\Gamma) \quad := \quad \frac{1}{\#(\mathbb{G})}\sum_{\mathsf{g}\in\mathbb{G}} \mathsf{g}.$$

We say Γ is *centred* if $|\mathsf{centre}(\Gamma)| < 1$. For any prime $p \in \mathbb{N}$, let

$$K_p \quad := \quad \min\left\{\frac{1}{2},\ \frac{4p-7}{4p+4}\right\}. \quad \text{Thus, } K_2 \;=\; \frac{1}{12}, \quad K_3 \;=\; \frac{5}{16}, \quad \text{and} \quad K_p \;=\; \frac{1}{2}, \ \text{for } p\geq 5.$$

Let $\mathcal{A} := (\mathbb{Z}_{/p})^s$ (where p is prime and $s\in\mathbb{N}$). If $\Phi : \mathcal{A}^{\mathbb{M}}\longrightarrow\mathcal{A}^{\mathbb{M}}$ is an LCA, then we say Φ is *bipartite* if $\Phi \;=\; 1+\Gamma\circ\sigma^{\mathsf{f}}$, where Γ is centred and $\mathsf{diam}[\Gamma] \;\leq\; K_p\cdot|\mathsf{f}|$. For example:

$\Phi \;=\; 1+\sigma^{\mathsf{f}}$ is bipartite for any nonzero $\mathsf{f}\in\mathbb{M}$ and any prime $p\in\mathbb{N}$.
$\Phi \;=\; 1+\sigma^{12}+\sigma^{13} \;=\; 1+(1+\sigma)\circ\sigma^{12}$ is bipartite for any prime $p\in\mathbb{N}$.
$\Phi \;=\; 1+\sigma^{14}+\sigma^{19} \;=\; 1+\left(\sigma^{-2}+\sigma^{3}\right)\circ\sigma^{16}$ is bipartite for any prime $p\geq 3$.
$\Phi \;=\; 1+\sigma^{2}+\sigma^{3} \;=\; 1+(1+\sigma)\circ\sigma^{2}$ is bipartite for any prime $p\geq 5$.

Our goal in this section is to prove:

Theorem 4.1. *Let $\mathcal{A} = (\mathbb{Z}_{/p})^s$, where p prime and $s\in\mathbb{N}$. If Φ is bipartite, then Φ is dispersive.* □

For any $N\in\mathbb{N}$, let $\left[N^{(i)}|_{i=0}^{\infty}\right]$ denote the *p-ary expansion* of N, so that $N \;=\; \sum_{i=0}^{\infty} N^{(i)}p^i$.

Let $\mathbb{L}(N) \;:=\; \left\{n\in[0...N]\ ;\ n^{(i)}\leq N^{(i)}, \text{for all } i\in\mathbb{N}\right\}$.

Lemma 4.2. (Lucas's Theorem)

(a) $\displaystyle\begin{bmatrix}N\\ n\end{bmatrix}_p \;=\; \prod_{i=0}^{\infty}\begin{bmatrix}N^{(i)}\\ n^{(i)}\end{bmatrix}_p$, *where we define* $\begin{bmatrix}N^{(i)}\\ n^{(i)}\end{bmatrix}_p := 0$ *if* $n^{(i)} > N^{(i)}$, *and* $\begin{bmatrix}0\\ 0\end{bmatrix}_p := 1$.

(b) *Thus,* $\begin{bmatrix}N\\ n\end{bmatrix}_p \neq\ 0$ *iff* $n\in\mathbb{L}(N)$. □

For example, suppose $\mathbb{M} = \mathbb{Z}$ and $\Phi = 1+\sigma$. If we interpret equation (5) in the light of Lemma 4.2, we get: $\displaystyle\Phi^N \;=\; \sum_{n\in\mathbb{L}(N)}\begin{bmatrix}N\\ n\end{bmatrix}_p\sigma^n.$

Lemma 4.3. *Let $r,\ H\ \in\ \mathbb{N}$.*

(a) *If $M < p^r$, and $N = M + p^r\cdot H$, then $\mathbb{L}(N) \;=\; \mathbb{L}(M) + p^r\cdot\mathbb{L}(H)$ (see Figure 1).*



FIGURE 1. Lemma 4.3.

(b) *If* $m \in \mathbb{L}(M)$, $h \in \mathbb{L}(H)$, *and* $n = m + p^r \cdot h$, *then* $\begin{bmatrix} N \\ n \end{bmatrix}_p = \begin{bmatrix} M \\ m \end{bmatrix}_p \cdot \begin{bmatrix} H \\ h \end{bmatrix}_p$. □

For example, suppose $p = 2$ and $N = 53 = 5 + 48 = 5 + 2^4 \cdot 3$. Then $M = 5$, $r = 4$, and $H = 3$, and

$$\begin{aligned} \mathbb{L}(53) &= \mathbb{L}(5) + 2^4 \cdot \mathbb{L}(3) = \{0, 1,\ 4, 5\} + 16 \cdot \{0, 1, 2, 3\} \\ &= \{0, 1,\ 4, 5,\quad 16, 17,\ 20, 21,\quad 32, 33,\ 37, 38,\quad 48, 49,\ 52, 53\}. \end{aligned}$$

If $\boldsymbol{\chi} = \bigotimes_{\mathsf{k} \in \mathbb{K}} \chi_{\mathsf{k}}$ is a character, then define $\mathsf{diam}[\boldsymbol{\chi}] := \max\{|\mathsf{k} - \mathsf{j}|\ ;\ \mathsf{k}, \mathsf{j} \in \mathbb{K}\}$. It follows:

LEMMA 4.4. *Let* Φ *be an LCA, and let* $S > 0$.

(a) *If* $\boldsymbol{\chi}$ *is a character, and* $S_0 = S + \mathsf{diam}[\boldsymbol{\chi}]$, *then* $\mathsf{rank}_S(\boldsymbol{\chi} \circ \Phi) \geq \mathsf{rank}_{S_0}(\Phi)$.

(b) *If* Γ *is an LCA, and* $S_0 = S + \mathsf{diam}[\Gamma]$, *then* $\mathsf{rank}_S(\Gamma \circ \Phi) \geq \mathsf{rank}_{S_0}(\Phi)$. □

COROLLARY 4.5. Φ *is dispersive if and only if, for any* $S_0 > 0$, *there is a subset* $\mathbb{J} \subset \mathbb{N}$ *of density 1 such that* $\lim_{\mathbb{J} \ni j \to \infty} \mathsf{rank}_{S_0}(\Phi^j) = \infty$. □

To prove Theorem 4.1, we'll use Lemma 4.3 to verify the condition of Corollary 4.5. For any $S_0 > 0$, define

$$\mathbb{J}(S_0) := \left\{N \in \mathbb{N}\ ;\ N = M_N + p^{r_N} H_N,\ \text{for some } H_N, r_N > 0 \text{ such that } M_N,\ S_0 < p^{r_N - 1}\right\}.$$

For example, if $p = 2$ and $S_0 = 7$, then $53 \in \mathbb{J}(7)$, because $53 = 5 + 2^4 \cdot 3$, so that $M_{53} = 5$, $r_{53} = 4$, and $H_{53} = 3$. Thus, $2^{r_{53}-1} = 2^3 = 8$, and $7 < 8$ and $5 < 8$. Note that $53 = 2^0 + 2^2 + 2^4 + 2^5$; thus, $53^{(3)} = 0$. This is exactly why $53 \in \mathbb{J}(7)$:

LEMMA 4.6. $\mathbb{J}(S_0) = \left\{N \in \mathbb{N}\ ;\ N \geq p \cdot S_0,\ \textit{and}\ N^{(r)} = 0 \textit{ for some } r \in \left(\log_p(S_0) \ldots \log_p(N)\right]\right\}$.

Proof. Suppose $N = M_N + p^{r_N} H_N$, for some $H_N, r_N > 0$ and $M_N \geq 0$, such that $M_N,\ S_0 < p^{r_N - 1}$. Let $r := r_N - 1$; then $N^{(r)} = 0$ and $\log_p(S_0) < r < \log_p(N)$.

Conversely, suppose $N^{(r)} = 0$, where $\log_p(S_0) < r < \log_p(N)$. Let $r_N := r + 1$; then $S_0 < p^r = p^{r_N - 1}$. Let $M_N := \sum_{i=0}^{r-1} N^{(i)} p^i$; then $M_N < p^r = p^{r_N - 1}$ also. Now let $H_N := \sum_{i=r_N}^{\infty} N^{(i)} p^{i - r_N}$; then $N = M_N + p^{r_N} H_N$. □

Lemma 4.7. $\mathsf{density}\left(\mathbb{J}(S_0)\right) = 1$.

Proof. Let $\mathbb{I} := [pS_0 \ldots \infty]$. Then $\mathbb{I}$ is a set of density one, and Lemma 4.6 implies that

$$\mathbb{I} \setminus \mathbb{J}(S_0) \quad = \quad \Big\{ N \in \mathbb{I}\ ;\ N^{(r)} \neq 0 \text{ for all } r \in \big(\log_p(S_0) \ldots \log_p(N)\big] \Big\},$$

which is a set of density zero. It follows that $\mathsf{density}\left(\mathbb{J}(S_0)\right) = \mathsf{density}\left(\mathbb{I}\right) = 1$. □

Lemma 4.8. *If $N \in \mathbb{J}(S_0)$, and $N = M + p^r H$, then $\Phi^N \ = \ \Phi^M \circ \Theta^H$, where $\Theta = \Phi^{(p^r)}$.*

Proof. Recall that $\Phi \ = \ 1 + \Gamma \circ \sigma^{\mathsf{f}}$. Thus,

$$\begin{aligned}
\Phi^N \ &\underset{(\mathrm{L})}{=} \ \sum_{n \in \mathbb{L}(N)} \begin{bmatrix} N \\ n \end{bmatrix}_p \Big(\Gamma \circ \sigma^{\mathsf{f}}\Big)^n \ \underset{(\ddagger)}{=} \ \sum_{m \in \mathbb{L}(M)} \ \sum_{h \in \mathbb{L}(H)} \begin{bmatrix} H \\ h \end{bmatrix}_p \begin{bmatrix} M \\ m \end{bmatrix}_p \Big(\Gamma \circ \sigma^{\mathsf{f}}\Big)^{(m + p^r h)} \\
&= \ \sum_{h \in \mathbb{L}(H)} \begin{bmatrix} H \\ h \end{bmatrix}_p \left(\sum_{m \in \mathbb{L}(M)} \begin{bmatrix} M \\ m \end{bmatrix}_p \left(\Gamma \circ \sigma^{\mathsf{f}}\right)^m \right) \circ \Big(\Gamma \circ \sigma^{\mathsf{f}}\Big)^{hp^r} \\
&\underset{(\dagger)}{=} \ \sum_{h \in \mathbb{L}(H)} \begin{bmatrix} H \\ h \end{bmatrix}_p \Phi^M \circ \Big(\Gamma \circ \sigma^{\mathsf{f}}\Big)^{p^r h} \ \underset{(\star)}{=} \ \Phi^M \circ \Theta^H.
\end{aligned}$$

(L) is by Lucas Theorem and ($\ddagger$) is by Lemma 4.3(b). ($\dagger$) is because $\Phi^M \ = \ \sum_{m \in \mathbb{L}(M)} \begin{bmatrix} M \\ m \end{bmatrix}_p \left(\Gamma \circ \sigma^{\mathsf{f}}\right)^m$. Finally, ($\star$) is because $\Theta \ = \ \left(1 + \Gamma \circ \sigma^{\mathsf{f}}\right)^{p^r} \underset{(\mathrm{L})}{=} 1 + (\Gamma \circ \sigma^{\mathsf{f}})^{p^r}$. Thus, $\Theta^H \underset{(\mathrm{L})}{=} \sum_{h \in \mathbb{L}(H)} \begin{bmatrix} H \\ h \end{bmatrix}_p \Big(\Gamma \circ \sigma^{\mathsf{f}}\Big)^{p^r h}$. □

Proof of Theorem 4.1. It suffices to verify the condition of Corollary 4.5. So, let $S_1 := S_0 + \mathsf{diam}\left[\Phi^M\right]$. Then

$$\mathsf{rank}_{S_0}\left(\Phi^N\right) \ \underset{(*)}{=} \ \mathsf{rank}_{S_0}\left(\Phi^M \circ \Theta^H\right) \ \underset{(\dagger)}{\geq} \ \mathsf{rank}_{S_1}\left(\Theta^H\right). \tag{6}$$

where ($*$) is by Lemma 4.8 and ($\dagger$) is by Lemma 4.4(b). Thus, we want to show that $\mathsf{rank}_{S_1}\left(\Theta^H\right) \xrightarrow[H \to \infty]{} \infty$ for H in a set of density 1. To do this, we'll use *gaps* in $\mathbb{L}\left(H\right)$. If $h_0, h_1 \in \mathbb{L}\left(H\right)$, we say that h_0 and h_1 ***bracket a gap*** if:

$$\textbf{(i)} \ \ h_1 \geq p \cdot h_0 \qquad \text{and} \qquad \textbf{(ii)} \ \ [h_0 ... h_1) \cap \mathbb{L}\left(H\right) \ = \ \emptyset.$$

Claim 1: *Let $h_0, h_1 \in \mathbb{L}\left(H\right)$, with $p \leq h_0 < h_1$, and suppose h_0 and h_1 bracket a gap in $\mathbb{L}\left(H\right)$. Then $\Big(\Gamma \circ \sigma^{\mathsf{f}}\Big)^{p^r h_0}$ and $\Big(\Gamma \circ \sigma^{\mathsf{f}}\Big)^{p^r h_1}$ are S_1-separated.*

Proof. Suppose $|h_0 - h_1| \ = \ w$. Then $\left(\sigma^{\mathsf{f}}\right)^{p^r h_0}$ and $\left(\sigma^{\mathsf{f}}\right)^{p^r h_1}$. are $(p^r \cdot w \cdot |\mathsf{f}|)$-separated. Thus, if $D = \mathsf{diam}\left[\Gamma\right]$, then $\Big(\Gamma \circ \sigma^{\mathsf{f}}\Big)^{p^r h_0}$ and $\Big(\Gamma \circ \sigma^{\mathsf{f}}\Big)^{p^r h_1}$ are W-separated, where

$$\begin{aligned}
W \ \ &:= \ \ p^r w |\mathsf{f}| \ - \ \left(\mathsf{diam}\left[\Gamma^{p_r h_0}\right] \ + \ \mathsf{diam}\left[\Gamma^{p_r h_1}\right]\right) \ \ = \ \ p^r w |\mathsf{f}| \ - \ (p^r h_0 D \ + \ p^r h_1 D) \\
&\geq \ \ p^r \cdot \Big(w|\mathsf{f}| - D \cdot (h_1 + h_0)\Big).
\end{aligned} \tag{7}$$



FIGURE 2. Claim 1 of Theorem 4.1.

(see Figure 2). We want $W \geq S_1$, or, equivalently, $W - \mathsf{diam}\left[\Phi^M\right] \geq S_0$ (because $S_1 = S_0 + \mathsf{diam}\left[\Phi^M\right]$). First, note that

$$\begin{aligned}\mathsf{diam}\left[\Phi^M\right] \quad &\leq \quad M \cdot |\mathsf{f}| \ + \ 2 \cdot \max_{m \in \mathbb{L}(M)} \mathsf{diam}\left[\Gamma^m\right] \quad = \quad M \cdot |\mathsf{f}| \ + \ 2M \cdot D \\ &= \quad M \cdot \Big(|\mathsf{f}| \ + \ 2D\Big) \quad \leq \quad p^{r-1} \cdot \Big(|\mathsf{f}| \ + \ 2D\Big). \end{aligned} \tag{8}$$

Thus,

$$\begin{aligned} W - \mathsf{diam}\left[\Phi^M\right] \quad &\underset{(*)}{\geq} \quad p^r \cdot \Big(w \cdot |\mathsf{f}| - D \cdot (h_1 + h_0)\Big) \ - \ p^{r-1} \cdot \Big(|\mathsf{f}| \ + \ 2D\Big) \\ &= \quad p^{r-1} \cdot \Big(pw \cdot |\mathsf{f}| \ - \ pD \cdot (h_1 + h_0) \ - \ |\mathsf{f}| \ - \ 2D\Big) \\ &\underset{(\dagger)}{\geq} \quad S_0 \cdot \Big(pw \cdot |\mathsf{f}| \ - \ pD \cdot (h_1 + h_0) \ - \ |\mathsf{f}| \ - \ 2D\Big). \end{aligned}$$

where $(*)$ is by equations (7) and (8), and $(\dagger)$ is because $S_0 < p^{r-1}$.

Thus, it suffices to show that

$$pw \cdot |\mathsf{f}| \ - \ pD \cdot (h_1 + h_0) \ - \ |\mathsf{f}| \ - \ 2D \quad \geq \quad 1.$$

To see this, observe that

$$\begin{aligned} &pw \cdot |\mathsf{f}| \ - \ pD \cdot (h_1 + h_0) \ - \ |\mathsf{f}| \ - \ 2D \\ &= \quad (pw - 1) \cdot |\mathsf{f}| - \Big[p \cdot (h_1 + h_0) - 2\Big] \cdot D \quad \underset{(\flat)}{\geq} \quad (pw - 1) \cdot |\mathsf{f}| - \Big[p \cdot (h_1 + h_0) - 2\Big] \cdot K_p \cdot |\mathsf{f}| \\ &= \quad \Big(pw - 1 - [p \cdot (h_1 + h_0) - 2]\, K_p\Big) \cdot |\mathsf{f}| \quad \underset{(*)}{\geq} \quad p \cdot (h_1 - h_0) - 1 - [p \cdot (h_1 + h_0) - 2]\, K_p \\ &= \quad p \cdot \Big((1 - K_p) \cdot h_1 - (1 + K_p) \cdot h_0\Big) \ - \ (1 + 2 \cdot K_p) \\ &\underset{(\dagger)}{\geq} \quad p \cdot \Big((1 - K_p) \cdot p - (1 + K_p)\Big) \cdot h_0 \ - \ 2 \quad \underset{(\ddagger)}{\geq} \quad p^2 \cdot \Big((1 - K_p) \cdot p - (1 + K_p)\Big) \ - \ 2 \\ &\underset{(\star)}{\geq} \quad \frac{3}{4} p^2 \ - \ 2 \quad \underset{(\diamond)}{\geq} \quad 3 - 2 \quad = \quad 1. \end{aligned}$$

($\flat$) is by hypothesis that Γ is bipartite. ($*$) is because $|\mathsf{f}| \geq 1$, and $w = h_1 - h_0$.

($\dagger$) is because $h_1 \geq p \cdot h_0$, and $K_p \leq \frac{1}{2}$. ($\ddagger$) is because $h_0 \geq p$.

($\star$) is because $K_p \leq \frac{4p-7}{4p+4} = \frac{p-\frac{7}{4}}{p+1}$, thus, $(p+1)K_p \leq p - \frac{7}{4} = p - 1 - \frac{3}{4}$; thus, $\frac{3}{4} \leq (p-1) - (p+1)K_p = (1-K_p)p - (1+K_p)$.

($\diamond$) is because $p \geq 2$, so $p^2 \geq 4$.

It follows that $W - \mathsf{diam}\left[\Phi^M\right] \geq S_0$, so that $W \geq S_1$. $\diamondsuit$ `Claim 1`

Let $\mathsf{rank}[H] := \#$ of gaps in $\mathbb{L}(H)$. Then Claim 1 implies that

$$\mathsf{rank}_{S_1}\left(\Theta^H\right) \quad \geq \quad \mathsf{rank}[H]. \tag{9}$$

Thus, we want to show that the number of gaps is large.

Suppose $i < k$. We say that i and k *bracket a zero-block* in the p-ary expansion of H if $H^{(i-1)} \neq 0 \neq H^{(k)}$, but $H^{(j)} = 0$, for all $i \leq j < k$. For example, suppose $p = 2$ and $H = 19$. Then 3 and 5 bracket a zero block in the binary expansion ...010011.

Claim 2: *If i and k bracket a zero-block in the p-ary expansion of H, then p^i and p^j bracket a gap in $\mathbb{L}(H)$.*

Proof. $H^{(i)} = 0$, so the largest element in $\mathbb{L}(H)$ less than p^i is

$$h_0 \quad = \quad \sum_{j=1}^{i-1} H^{(j)} \cdot p^j \quad \leq \quad \sum_{j=1}^{i-1} (p-1) \cdot p^j \quad = \quad p^i - 1.$$

Now, $k = \min\left\{j > i\,;\ H^{(j)} \neq 0\right\}$, so $h_1 = p^k$ is the smallest element in $\mathbb{L}(H)$ greater than p^i. Also, $h_1 \geq p^{i+1} > p \cdot (p^i - 1) \geq p \cdot h_0$. $\diamondsuit$ `Claim 2`

Let $\#\mathbf{ZB}(H) := \#$of zero-blocks in the p-ary expansion of H.

Then Claim 2 implies that

$$\mathsf{rank}[H] \quad \geq \quad \#\mathbf{ZB}(H). \tag{10}$$

Define $\mathbb{H} := \left\{H \in \mathbb{N}\,;\ \#\mathbf{ZB}(H) \geq \frac{1}{p^3}\log_p(H)\right\}$.

Claim 3: $\mathsf{density}(\mathbb{H}) = 1$.

Proof. Observe that $\#\mathbf{ZB}(H)$ is no less than the number of occurrences of the word "101" in the p-ary expansion of H (because 101 is a zero-block). Let

$$\mathbb{H}' \quad := \quad \left\{H \in \mathbb{N}\,;\ (\#\text{ of occurrences of “101”}) \geq \frac{1}{p^3}\log_p(H)\right\}.$$

Then $\mathbb{H}' \subset \mathbb{H}$. The Weak Law of Large Numbers implies $\mathsf{density}(\mathbb{H}') = 1$. $\diamondsuit$ `Claim 3`

Define $\mathbb{J} := \left\{N \in \mathbb{J}(S_0)\,;\ N = M_N + p^{r_N} H_N,\ \text{where } r_N \leq \frac{1}{2}\log_p(N),\ \text{and } H_N \in \mathbb{H}\right\}$.

Claim 4: $\mathsf{density}(\mathbb{J}) = 1$.

Proof. $\mathbb{J} = \mathbb{J}_1 \cap \mathbb{J}_2$, where

$$\begin{aligned} \mathbb{J}_1 &:= \{N \in \mathbb{J}(S_0) \ ; \ N = M_N + p^{r_N} H_N, \text{ where } H_N \in \mathbb{H}\} \\ \text{and } \ \mathbb{J}_2 &:= \left\{N \in \mathbb{J}(S_0) \ ; \ N = M_N + p^{r_N} H_N, \text{ where } r_N \leq \frac{1}{2}\log_p(N)\right\}. \end{aligned}$$

Now, $\mathsf{density}(\mathbb{J}_1) = 1$ by Lemma 4.7 and Claim 3. To see that $\mathsf{density}(\mathbb{J}_2) = 1$, note that

$$\mathbb{J}(S_0) \setminus \mathbb{J}_2 \quad \subset \quad \left\{N \in \mathbb{N} \ ; \ N^{(r)} \neq 0 \text{ for all } r \in \left(\log_p(S_0) \ldots \frac{1}{2}\log_p(N)\right]\right\},$$

which is a set of density zero. ◇ `Claim 4`

If $N = M_N + p^{r_N} H_N$ is an element of $\mathbb{J}$, then

$$\log_p(H_N) \quad \geq \quad \log_p(N) - r_N \quad \geq \quad \log_p(N) - \frac{1}{2}\log_p(N) \quad = \quad \frac{1}{2}\log_p(N). \tag{11}$$

Thus,

$$\begin{aligned} \mathsf{rank}_{S_0}\left(\Phi^N\right) \ &\underset{(\heartsuit)}{\geq} \ \mathsf{rank}_{S_1}\left(\Theta^{H_N}\right) \ \underset{(\diamondsuit)}{\geq} \ \mathsf{rank}\left[H_N\right] \ \underset{(\clubsuit)}{\geq} \ \#\mathbf{ZB}\left(H_N\right) \\ &\underset{(*)}{\geq} \ \frac{1}{p^3}\log_p(H_N) \ \underset{(\spadesuit)}{\geq} \ \frac{1}{2p^3}\log_p(N). \end{aligned}$$

Here, $(\heartsuit)$ is by equation (6), $(\diamondsuit)$ is by equation (9), $(\clubsuit)$ is by equation (10), $(\spadesuit)$ is by equation (11), and $(*)$ is because $H \in \mathbb{H}$ by hypothesis.

Thus $\displaystyle\lim_{\mathbb{J} \ni N \to \infty} \mathsf{rank}_{S_0}\left(\Phi^N\right) \quad \geq \quad \frac{1}{2p^3} \lim_{\mathbb{J} \ni N \to \infty} \log_p(N) \ = \ \infty.$ □

5. *Uniform Mixing and Dispersion Mixing*

A measure $\mu \in \mathcal{M}(\mathcal{A}^{\mathbb{Z}})$ is ***uniformly mixing*** if, for any $\epsilon > 0$, there is some $M > 0$ such that, for any cylinder subsets $\mathfrak{L} \subset \mathcal{A}^{(-\infty \ldots 0]}$ and $\mathfrak{R} \subset \mathcal{A}^{[0 \ldots \infty)}$, and any $m > M$,

$$\mu\left[\sigma^m(\mathfrak{L}) \ \cap \mathfrak{R}\right] \quad \underset{\epsilon}{\sim} \quad \mu\left[\mathfrak{L}\right] \cdot \mu\left[\mathfrak{R}\right] \tag{12}$$

(here, "$x \underset{\epsilon}{\sim} y$" means $|x - y| < \epsilon$.)

EXAMPLE 5.1:

⟨a⟩ Any mixing N-step Markov chain is uniformly mixing. (See §6).

⟨b⟩ If $\nu \in \mathcal{M}(\mathcal{B}^{\mathbb{Z}})$ is uniformly mixing, and $\Psi : \mathcal{B}^{\mathbb{Z}} \longrightarrow \mathcal{A}^{\mathbb{Z}}$ is a block map, then $\mu := \Phi(\nu)$ is also uniformly mixing. (If Ψ has local map $\psi : \mathcal{B}^{[-\ell \ldots r]} \longrightarrow \mathcal{A}$, then replace the M in (12) with $M + \ell + r + 1$).

⟨c⟩ Hence, if $\mathfrak{F} \subset \mathcal{B}^{\mathbb{Z}}$ is an SFT, and $\mathfrak{S} := \Psi(\mathfrak{F}) \subset \mathcal{A}^{\mathbb{Z}}$ a sofic shift, and $\nu \in \mathcal{M}(\mathfrak{F})$ is any mixing N-step Markov chain, then $\mu := \Phi(\nu)$ is a uniformly mixing measure on $\mathfrak{S}$. We call μ a ***quasi-Markov measure***. ◇

We say that μ is ***harmonically bounded*** (HB) if there is some $C < 1$ such that $|\langle \boldsymbol{\chi}, \mu \rangle| < C$ for all $\boldsymbol{\chi} \in \widehat{\mathcal{A}^{\mathbb{Z}}}$ except $\boldsymbol{\chi} = \mathbb{1}$. The goal of this section is to prove:

THEOREM 5.2. *Let $\mathcal{A}$ be a finite abelian group. If $\mu \in \mathcal{M}(\mathcal{A}^{\mathbb{Z}})$ is uniformly mixing and harmonically bounded, then μ is dispersion mixing.* □

We will then apply Theorem 5.2 to get:

COROLLARY 5.3. *Let $\mathcal{A} = \mathbb{Z}_{/p}$, where p is prime. If $\mu \in \mathcal{M}(\mathcal{A}^{\mathbb{Z}})$ is a mixing quasi-Markov measure, then μ is asymptotically randomized by any dispersive LCA.* □

Harmonic boundedness and entropy:

LEMMA 5.4. *Let $\mathcal{A} = (\mathbb{Z}_{/p})^s$, where p is prime and $s \in \mathbb{N}$. If $\mu \in \mathcal{M}(\mathcal{A}^{\mathbb{Z}})$ and $h(\mu, \sigma) > (s-1) \cdot \log_2(p)$, then μ is harmonically bounded.*

Proof. Suppose μ was not HB. Then for any $\alpha > 0$, we can find $\mathbb{1} \neq \boldsymbol{\chi} \in \widehat{\mathcal{A}^{\mathbb{Z}}}$ with $|\langle \boldsymbol{\chi}, \mu \rangle| > 1 - \alpha$. Let $\mathcal{I} := \mathrm{image}\,(\boldsymbol{\chi}) \subset \mathbb{T}^1$, and let $\nu := \boldsymbol{\chi}(\mu) \in \mathcal{M}(\mathcal{I})$ be the projected measure on $\mathcal{I}$. Thus, $\langle \boldsymbol{\chi}, \mu \rangle = \sum_{i \in \mathcal{I}} i \cdot \nu\{i\}$. The following four claims are easy to check.

CLAIM 1: *For any $\beta > 0$, there exists $\alpha > 0$ such that, for any probability measure $\nu \in \mathcal{M}(\mathcal{I})$ with $\left| \sum_{i \in \mathcal{I}} i \cdot \nu\{i\} \right| > 1 - \alpha$, there is some $i_0 \in \mathcal{I}$ with $\nu\{i_0\} > 1 - \beta$.* ◇

Suppose $\boldsymbol{\chi} = \bigotimes_{k \in \mathbb{K}} \chi_k$, where $\mathbb{K} \subset [0...K]$ and $K \in \mathbb{K}$. Thus, if $\boldsymbol{\xi} := \bigotimes_{k \in \mathbb{K} \setminus \{K\}} \chi_k$, then $\boldsymbol{\chi} = \boldsymbol{\xi} \otimes \chi_K$. For any $\mathbf{b} \in \mathcal{A}^{[0...K)}$, let $\mu_K^{(\mathbf{b})}$ be the conditional measure on the Kth coordinate, and let $\nu_K^{(\mathbf{b})} := \chi_K\left(\mu_K^{(\mathbf{b})}\right) \in \mathcal{M}(\mathcal{I})$ be the projected measure on $\mathcal{I}$.

CLAIM 2: *For any $\gamma > 0$, there exists $\beta > 0$ such that, if $\exists\, i_0 \in \mathcal{I}$ with $\nu\{i_0\} > 1 - \beta$, then there is a subset $\mathfrak{B} \subset \mathcal{A}^{[0...K)}$ with $\mu[\mathfrak{B}] > 1 - \gamma$, such that, for every $\mathbf{b} \in \mathfrak{B}$, there is some $i_{\mathbf{b}} \in \mathcal{I}$ with $\nu_K^{(\mathbf{b})}\{i_{\mathbf{b}}\} > 1 - \gamma$. Thus, if $\mathcal{P}_{\mathbf{b}} = \chi_K^{-1}\{i_{\mathbf{b}}\} \subset \mathcal{A}$, then $\mu_K^{(\mathbf{b})}[\mathcal{P}_{\mathbf{b}}] > 1 - \gamma$. (Observe that $\#(\mathcal{P}_{\mathbf{b}}) \leq p^{s-1}$ for all $\mathbf{b} \in \mathcal{A}^{[0...K)}$.)* ◇

For any measure $\rho \in \mathcal{M}(\mathcal{A})$, define $H(\rho) := -\sum_{a \in \mathcal{A}} \rho\{a\} \log_2\left(\rho\{a\}\right)$. Recall (e.g. [**Pet89**, Proposition 5.2.12]) that the σ-entropy of μ can be computed:

$$h(\mu, \sigma) \quad = \quad \lim_{N \to \infty} \sum_{\mathbf{b} \in \mathcal{A}^{[0...N)}} \mu[\mathbf{b}] \cdot H\left(\mu_N^{(\mathbf{b})}\right) \tag{13}$$

CLAIM 3: *For any $\delta > 0$, there exists $\gamma_1 > 0$ such that, for any probability measure ρ on $\mathcal{A}$, if there is a subset $\mathcal{P} \subset \mathcal{A}$ with $\#(\mathcal{P}) \leq p^{s-1}$ and $\rho[\mathcal{P}] > 1 - \gamma_1$, then $H(\rho) < (s-1) \cdot \log_2(p) + \delta$.* ◇

CLAIM 4: *For any $\epsilon > 0$, and $S > 0$, there exist $\delta, \gamma_2 > 0$ such that, for any $K \in \mathbb{N}$ and probability measure μ on $\mathcal{A}^{[0...K]}$, if there is a subset $\mathfrak{B} \subset \mathcal{A}^{[0...K)}$ with $\mu[\mathfrak{B}] > 1 - \gamma_2$, such that, for all $\mathbf{b} \in \mathfrak{B}$, $H\left(\mu_K^{(\mathbf{b})}\right) < S - \delta$, then $\sum_{\mathbf{b} \in \mathcal{A}^{[0...K)}} \mu[\mathbf{b}] \cdot H\left(\mu_K^{(\mathbf{b})}\right) < S - \epsilon$.* ◇

Now, set $S := (s-1) \cdot \log_2(p)$. For any $\epsilon > 0$, find $\delta, \gamma_2 > 0$ as in Claim 4. Then find $\gamma_1 > 0$ as in Claim 3, and let $\gamma := \min\{\gamma_1, \gamma_2\}$. Next, find β as in Claim 2 and then find α as in Claim 1. Finally, find $\boldsymbol{\chi} \in \widehat{\mathcal{A}^{\mathbb{Z}}}$ with $|\langle \boldsymbol{\chi}, \mu \rangle| > 1 - \alpha$. It then follows from

Claims 1-4 that $\displaystyle\sum_{\mathbf{b}\in\mathcal{A}^{[0...K)}} \mu\left[\mathbf{b}\right]\cdot H\left(\mu_N^{(\mathbf{b})}\right) \ < \ (s-1)\cdot\log_2(p)-\epsilon$. But the limit in (13) is a decreasing limit, so we conclude that $h(\mu,\sigma) \ < \ (s-1)\cdot\log_2(p)-\epsilon$. Since this is true for any $\epsilon>0$, we conclude that $h(\mu,\sigma) \ \leq \ (s-1)\cdot\log_2(p)$, contradicting our hypothesis. □

Corollary 5.5. *If* $\mathcal{A}=\mathbb{Z}_{/p}$ (where p is prime)*, and* $h(\mu,\sigma)>0$*, then* μ *is harmonically bounded.* □

Say μ is *uniformly multiply mixing* if, for any $\epsilon>0$, there is some $S>0$ such that, for any $R>0$, if $\mathbb{K}_0,\mathbb{K}_1,\ldots,\mathbb{K}_R\subset\mathbb{M}$ are finite, mutually S-separated subsets of $\mathbb{M}$, and $\mathfrak{U}_0\subset\mathcal{A}^{\mathbb{K}_0},\ldots,\mathfrak{U}_R\subset\mathcal{A}^{\mathbb{K}_R}$ are cylinder sets, then:

$$\mu\left(\bigcap_{r=0}^{R}\mathfrak{U}_r\right) \ \underset{R\epsilon}{\widetilde{\quad}} \ \prod_{r=0}^{R}\mu\left(\mathfrak{U}_r\right). \tag{14}$$

Lemma 5.6. *If* $\mu\in\mathcal{M}(\mathcal{A}^{\mathbb{Z}})$ *is uniformly mixing, then* μ *is uniformly multiply mixing.*

Proof. (*by induction on* R). The case $R=1$ is just uniform mixing. Suppose (14) is true for all $R'<R$. Find $S>0$ so that, if $\mathbb{K}_0,\ldots,\mathbb{K}_R$ are mutually S-separated, then

$$\mu\left(\bigcap_{r=0}^{R}\mathfrak{U}_r\right) \ = \ \mu\left(\mathfrak{U}_0\cap\bigcap_{r=1}^{R}\mathfrak{U}_r\right) \ \underset{\epsilon}{\sim} \ \mu\left(\mathfrak{U}_0\right)\cdot\mu\left(\bigcap_{r=1}^{R}\mathfrak{U}_r\right) \ \underset{(R-1)\epsilon}{\widetilde{\quad\quad}} \ \mu\left(\mathfrak{U}_0\right)\cdot\prod_{r=1}^{R}\mu\left(\mathfrak{U}_r\right),$$

where "$\underset{\epsilon}{\sim}$" comes by setting $R'=1$, and "$\underset{(R-1)\epsilon}{\widetilde{\quad\quad}}$" comes by setting $R'=R-1$. □

Lemma 5.7. *Suppose* $\mu\in\mathcal{M}(\mathcal{A}^{\mathbb{Z}})$ *is uniformly multiply mixing. For any* $\epsilon>0$ *and* $R\in\mathbb{N}$*, there is some* $S>0$ *such that: if* $\mathbb{K}_0,\ldots,\mathbb{K}_R\subset\mathbb{Z}$ *are* S*-separated sets, and, for all* $r\in[0...R]$*,* $\boldsymbol{\chi}_r:\mathcal{A}^{\mathbb{K}_r}\longrightarrow\mathbb{C}$ *are characters, and* $\boldsymbol{\chi}=\displaystyle\prod_{r=0}^{R}\boldsymbol{\chi}_r$*, then*

$$\langle\boldsymbol{\chi},\ \mu\rangle \ \underset{\epsilon/2}{\widetilde{\quad}} \ \prod_{r=0}^{R}\langle\boldsymbol{\chi}_r,\ \mu\rangle.$$ □

Proof of Theorem 5.2. Let $\epsilon>0$. We want to find $S>0$ and $R>0$ such that, if $\boldsymbol{\chi}$ is any character, and $\mathsf{rank}_S(\boldsymbol{\chi})>R$, then $|\langle\boldsymbol{\chi},\mu\rangle|<\epsilon$.

Let $C<1$ be the harmonic bound. Find $R\in\mathbb{N}$ such that $C^R<\epsilon/2$. Let $S>0$ be as in Lemma 5.7. Suppose $\mathsf{rank}_S(\boldsymbol{\chi})>R$, and let $\boldsymbol{\chi} \ := \ \displaystyle\bigotimes_{r=0}^{R}\boldsymbol{\chi}_r$, where $\boldsymbol{\chi}_r:\mathcal{A}^{\mathbb{K}_r}\longrightarrow\mathbb{C}$ are characters, and $\mathbb{K}_0,\ldots,\mathbb{K}_R\subset\mathbb{Z}$ are S-separated. Then Lemma 5.7 implies:

$$\langle\boldsymbol{\chi},\ \mu\rangle \ \underset{\epsilon/2}{\widetilde{\quad}} \ \prod_{r=0}^{R}\langle\boldsymbol{\chi}_r,\ \mu\rangle. \tag{15}$$

By harmonic boundedness, we know $|\langle\boldsymbol{\chi}_r,\ \mu\rangle|<C$ for all $r\in[0...R]$. Thus, (15) implies:

$$|\langle\boldsymbol{\chi},\ \mu\rangle| \ \underset{\epsilon/2}{\widetilde{\quad}} \ \prod_{r=0}^{R}|\langle\boldsymbol{\chi}_r,\ \mu\rangle| \ < \ \prod_{r=0}^{R}C \ = \ C^{R+1} \ < \ C^R \ < \ \epsilon/2.$$ □

Proof of Corollary 5.3. From examples 5.1(a) and 5.1(b), we know μ is uniformly mixing. Any mixing quasi-Markov measure has nonzero entropy, so Corollary 5.5 says that μ is harmonically bounded. Theorem 5.2 says μ is dispersion mixing. Theorem 3.1 says μ is asymptotically randomized by any dispersive CA. □

6. *Markov Words*

If $m, n \in \mathbb{Z}$, and $m \leq n$, let $\mathcal{A}^{[m...n)}$ be the set of all ***words*** of the form $\mathbf{a} = [a_m,\ a_{m+1}, \ldots, a_{n-1}]$. Let $\mathcal{A}^* := \bigcup_{-\infty<m<n<\infty} \mathcal{A}^{[m...n)}$ be the set of all finite words. Elements of $\mathcal{A}^*$ are denoted by boldfaced letters (e.g. $\mathbf{a}$, $\mathbf{b}$, $\mathbf{c}$), and subsets by gothic letters (e.g. $\mathfrak{A}$, $\mathfrak{B}$, $\mathfrak{C}$). Concatenation of words is indicated by juxtaposition. Thus, if $\mathbf{a} = [a_0 \ldots a_n]$ and $\mathbf{b} = [b_0 \ldots b_m]$, then $\mathbf{ab} = [a_0 \ldots a_n b_0 \ldots b_m]$.

If $V > 0$ and $\mathbf{v} \in \mathcal{A}^{[-V...V)}$, we say that $\mathbf{v}$ is a ***Markov word*** for μ if (in the terminology of §1), $\mathbf{v}$ isolates $(-\infty... -V)$ from $[V...\infty)$.

EXAMPLE 6.1:

⟨a⟩ If μ is an N-step Markov shift, and $N \leq 2V$, then every $\mathbf{v} \in \mathcal{A}^{[-V...V)}$ is a Markov word.

⟨b⟩ Let $\mathfrak{F} \subset \mathcal{B}^{\mathbb{Z}}$ be a subshift of finite type, let $\Psi : \mathfrak{F} \longrightarrow \mathcal{A}^{\mathbb{Z}}$ be a block map, so that $\mathfrak{S} := \Psi(\mathfrak{F})$ is a sofic shift. Let ν be a Markov measure on $\mathfrak{F}$ and let $\mu := \Psi(\nu)$. If $s \in \mathfrak{S}_{[-V...V]}$ is a synchronizing word for Ψ, then s is a Markov word for μ. ♢

PROPOSITION 6.2. *If $\mu \in \mathcal{M}(\mathcal{A}^{\mathbb{Z}})$ is mixing and has a Markov word, then μ is uniformly mixing.*

Proof. Fix $\epsilon > 0$. For any words $\mathbf{a}, \mathbf{b} \in \mathcal{A}^*$, the mixing of μ implies that there is some $M_\epsilon(\mathbf{a}, \mathbf{b}) < \infty$ such that, for all $m > M_\epsilon(\mathbf{a}, \mathbf{b})$, $\mu\left(\sigma^m[\mathbf{a}] \cap [\mathbf{b}]\right) \underset{\epsilon}{\sim} \mu[\mathbf{a}] \cdot \mu[\mathbf{b}]$. Our goal is to find some $M > 0$ such that $M_\epsilon(\mathbf{a}, \mathbf{b}) < M$ for all $\mathbf{a}, \mathbf{b} \in \mathcal{A}^*$.

Let $\mathbf{v} \in \mathcal{A}^*$ be a Markov word for μ.

CLAIM 1: *Let $\mathbf{u}, \mathbf{w}, \mathbf{u}', \mathbf{w}' \in \mathcal{A}^*$, and consider the words $\mathbf{uvw}$ and $\mathbf{u}'\mathbf{v}\mathbf{w}'$. We have: $M_\epsilon(\mathbf{uvw},\ \mathbf{u}'\mathbf{v}\mathbf{w}') = M_\epsilon(\mathbf{vw},\ \mathbf{u}'\mathbf{v})$.*

Proof. Define transition probabilities: $\mu(\mathbf{u} \dashleftarrow \mathbf{v}) := \mu(\mathbf{uv})/\mu(\mathbf{v})$ and $\mu(\mathbf{v} \dashrightarrow \mathbf{w}) := \mu(\mathbf{vw})/\mu(\mathbf{v})$. If $m > M_\epsilon(\mathbf{vw},\ \mathbf{u}'\mathbf{v})$, then

$$\begin{aligned} \mu\left(\sigma^m[\mathbf{uvw}] \cap [\mathbf{u}'\mathbf{v}\mathbf{w}']\right) &= \mu(\mathbf{u} \dashleftarrow \mathbf{v}) \cdot \mu\left(\sigma^m[\mathbf{vw}] \cap [\mathbf{u}'\mathbf{v}]\right) \cdot \mu(\mathbf{v} \dashrightarrow \mathbf{w}') && (16)\\ &\underset{\epsilon}{\sim} \mu(\mathbf{u} \dashleftarrow \mathbf{v}) \cdot \mu[\mathbf{vw}] \cdot \mu[\mathbf{u}'\mathbf{v}] \cdot \mu(\mathbf{v} \dashrightarrow \mathbf{w}') && (17)\\ &= \mu[\mathbf{uvw}] \cdot \mu[\mathbf{u}'\mathbf{v}\mathbf{w}']. && (18) \end{aligned}$$

(16) and (18) are because $\mathbf{v}$ is a Markov word; (17) is because $m > M_\epsilon(\mathbf{vw}, \mathbf{u}'\mathbf{v})$.
♢ `Claim 1`

If $\mathbf{a} \in \mathcal{A}^*$, we say that $\mathbf{v}$ *occurs* in $\mathbf{a}$ if $\mathbf{a}\big|_{[n-V\ldots n+V)} = \mathbf{v}$ for some n.

CLAIM 2: *There is some $N > 0$ such that $\mu\big\{\mathbf{a} \in \mathcal{A}^{[0\ldots N]} \ ; \ \mathbf{v}$ occurs in $\mathbf{a}\big\} > 1 - \epsilon$.*

Proof. By ergodicity, find N such that $\mu\left(\bigcup_{n=0}^{N} \sigma^n [\mathbf{v}]\right) > 1 - \epsilon$. ◇ `Claim 2`

Let $\mathcal{A}^*_{\mathbf{v}}$ be the set of words (of length at least N) in $\mathcal{A}^*$ with $\mathbf{v}$ occuring in the last $(N+V)$ coordinates, and let ${}_{\mathbf{v}}\mathcal{A}^*$ be the set of all words in $\mathcal{A}^*$ with $\mathbf{v}$ occuring in the first $(N+V)$ coordinates. Then Claim 2 implies that:

$$\mu(\mathcal{A}^*_{\mathbf{v}}) \quad > \quad 1-\epsilon \quad \text{and} \quad \mu({}_{\mathbf{v}}\mathcal{A}^*) \quad > \quad 1-\epsilon. \tag{19}$$

Let $\mathcal{A}^{<N} := \bigcup_{n=1}^{N} \mathcal{A}^{[0\ldots n]}$. Then

$$\begin{array}{rcl} \mathcal{A}^*_{\mathbf{v}} & = & \big\{\mathbf{uvw} \ ; \ \mathbf{u} \in \mathcal{A}^* \text{ and } \ \mathbf{w} \in \mathcal{A}^{<N}\big\}. \\ \text{and } \ {}_{\mathbf{v}}\mathcal{A}^* & = & \big\{\mathbf{u'vw'} \ ; \ \mathbf{u'} \in \mathcal{A}^{<N} \text{ and } \ \mathbf{w'} \in \mathcal{A}^*\big\}, \end{array} \tag{20}$$

$$\begin{aligned} \text{Define} \quad M_1 \ &:= \max_{\mathbf{a}\in\mathcal{A}^*_{\mathbf{v}}} \max_{\mathbf{b}\in{}_{\mathbf{v}}\mathcal{A}^*} M_\epsilon(\mathbf{a},\mathbf{b}) \underset{(*)}{=\!=} \max_{\substack{\mathbf{u}\in\mathcal{A}^* \\ \mathbf{w}\in\mathcal{A}^{<N}}} \max_{\substack{\mathbf{u'}\in\mathcal{A}^{<N} \\ \mathbf{w'}\in\mathcal{A}^*}} M_\epsilon\left(\mathbf{uvw}, \ \mathbf{u'vw'}\right) \\ &\underset{(\dagger)}{=\!=} \max_{\mathbf{w},\mathbf{u'}\in\mathcal{A}^{<N}} M_\epsilon\left(\mathbf{vw}, \ \mathbf{u'v}\right). \end{aligned}$$

where $(*)$ is by equation (20) and $(\dagger)$ is by Claim 1. Likewise, define

$$\begin{array}{rclcl} M_2 & := & \displaystyle\max_{\mathbf{a}\in\mathcal{A}^*_{\mathbf{v}}} \max_{\mathbf{b}\in\mathcal{A}^{<N}} M_\epsilon(\mathbf{a},\mathbf{b}) & = & \displaystyle\max_{\mathbf{w}\in\mathcal{A}^{<N}} \max_{\mathbf{b}\in\mathcal{A}^{<N}} M_\epsilon(\mathbf{vw}, \ \mathbf{b}), \\ M_3 & := & \displaystyle\max_{\mathbf{a}\in\mathcal{A}^{<N}} \max_{\mathbf{b}\in{}_{\mathbf{v}}\mathcal{A}^*} M_\epsilon(\mathbf{a},\mathbf{b}) & = & \displaystyle\max_{\mathbf{a}\in\mathcal{A}^{<N}} \max_{\mathbf{u'}\in\mathcal{A}^{<N}} M_\epsilon(\mathbf{a}, \ \mathbf{u'v}), \\ \text{and } \ M_4 & := & \displaystyle\max_{\mathbf{a}\in\mathcal{A}^{<N}} \max_{\mathbf{b}\in\mathcal{A}^{<N}} M_\epsilon(\mathbf{a},\mathbf{b}). & & \end{array}$$

Thus, $M_1, \ldots, M_4$ each maximizes a finite collection of finite values, so each is finite. Thus, $M := \max\{M_1, \ldots, M_4\}$ is finite.

CLAIM 3: *For any $\mathbf{a}, \mathbf{b} \in \mathcal{A}^*$, $\quad M_\epsilon(\mathbf{a},\mathbf{b}) < M$.*

Proof. If $\mathbf{a} \in \mathcal{A}^{<N} \cup \mathcal{A}^*_{\mathbf{v}}$ and $\mathbf{b} \in \mathcal{A}^{<N} \cup \ {}_{\mathbf{v}}\mathcal{A}^*$, then $M_\epsilon(\mathbf{a},\mathbf{b}) < M$ by definition.

So, suppose $\mathbf{a} \notin \mathcal{A}^{<N} \cup \mathcal{A}^*_{\mathbf{v}}$. Then equation (19) implies that $\mu[\mathbf{a}] < \epsilon$. Hence, for any $m \in \mathbb{N}$, $\mu(\sigma^m[\mathbf{a}] \cap \mathbf{b}) < \epsilon$ and $\mu[\mathbf{a}] \cdot \mu[\mathbf{b}] < \epsilon$. Thus, $\mu(\sigma^m[\mathbf{a}] \cap \mathbf{b}) \underset{\epsilon}{\sim} \mu[\mathbf{a}] \cdot \mu[\mathbf{b}]$ automatically. Hence, $M_\epsilon(\mathbf{a},\mathbf{b}) = 0 < M$.

Likewise, if $\mathbf{b} \notin \mathcal{A}^{<N} \cup \ {}_{\mathbf{v}}\mathcal{A}^*$, then $M_\epsilon(\mathbf{a},\mathbf{b}) = 0 < M$. ◇ `Claim 3`

Thus, μ is uniformly mixing. □

COROLLARY 6.3. *If μ is harmonically bounded, mixing and has a Markov word, then μ is asymptotically randomized by $\Phi = 1 + \sigma$.*

Proof. Combine Proposition 6.2 with Theorems 3.1 and 5.2. □

7. *Lucas Mixing*

Throughout this section, let $D := 1$, so that $\mathbb{M} = \mathbb{Z}$. Let $\mathcal{A} := (\mathbb{Z}_{/p})^s$, where $p \in \mathbb{N}$ is prime, and $s \in \mathbb{N}$. Let $\Phi := 1 + \sigma$. We will introduce a condition on μ which is weaker than dispersion mixing, and which is both sufficient and *necessary* for asymptotic randomization.

Let $\boldsymbol{\chi} \in \widehat{\mathcal{A}^{\mathbb{Z}}}$, and suppose $\boldsymbol{\chi} = \bigotimes_{\mathsf{k}\in\mathbb{K}} \chi_{\mathsf{k}}$. We define $[\![\boldsymbol{\chi}]\!] := \max(\mathbb{K}) - \min(\mathbb{K})$, and define

$$\langle\!\langle \boldsymbol{\chi} \rangle\!\rangle \quad := \quad p^r, \quad \text{where} \quad r := \left\lceil \log_p [\![\boldsymbol{\chi}]\!] \right\rceil .$$

It follows from Lucas' Theorem that $\Phi^{\langle\!\langle \boldsymbol{\chi} \rangle\!\rangle} = 1 + \sigma^{\langle\!\langle \boldsymbol{\chi} \rangle\!\rangle}$. Thus, for any $h \in \mathbb{N}$,

$$\Phi^{h\cdot\langle\!\langle \boldsymbol{\chi} \rangle\!\rangle} \quad = \quad \sum_{\ell \in \mathbb{L}(h)} \begin{bmatrix} h \\ \ell \end{bmatrix}_p \sigma^{\langle\!\langle \boldsymbol{\chi} \rangle\!\rangle\cdot\ell}, \quad \text{and thus,} \quad \boldsymbol{\chi}\circ\Phi^{h\cdot\langle\!\langle \boldsymbol{\chi} \rangle\!\rangle} \quad = \quad \bigotimes_{\ell \in \mathbb{L}(h)} \begin{bmatrix} h \\ \ell \end{bmatrix}_p \boldsymbol{\chi}\circ\sigma^{\langle\!\langle \boldsymbol{\chi} \rangle\!\rangle\cdot\ell}.$$

Observe that $\mathbb{K}+p^r\ell$ and $\mathbb{K}+p^r\ell'$ are disjoint for any $\ell \neq \ell' \in \mathbb{L}(h)$. Hence, if $L := \#(\mathbb{L}(h))$, then $\boldsymbol{\chi}\circ\Phi^{h\cdot\langle\!\langle \boldsymbol{\chi} \rangle\!\rangle}$ is a product of L 'disjoint translates' of $\boldsymbol{\chi}$.

If μ is a measure on $\mathcal{A}^{\mathbb{Z}}$, we say that μ is ***Lucas mixing*** if, for any nontrivial character $\boldsymbol{\chi} \in \widehat{\mathcal{A}^{\mathbb{Z}}}$, there is a subset $\mathbb{H} \subset \mathbb{N}$ of Cesàro density one such that $\lim_{\mathbb{H}\ni h\to\infty} \left\langle \boldsymbol{\chi}\circ\Phi^{h\cdot\langle\!\langle \boldsymbol{\chi} \rangle\!\rangle},\ \mu \right\rangle = 0$. Our goal in this section is to prove:

THEOREM 7.1. $\Big(\Phi = 1+\sigma$ *asymptotically randomizes* $\mu\Big) \iff \Big(\mu$ *is Lucas mixing*$\Big)$. □

It is relatively easy to see that:

LEMMA 7.2. *If μ is dispersion-mixing, then μ is Lucas mixing.* □

Thus, the "$\Longleftarrow$" direction of Theorem 7.1 is an extension of Theorem 3.1, in the case $\Phi = 1+\sigma$. The "$\Longrightarrow$" direction makes this the strongest possible extension for this LCA.

Set $S := [\![\boldsymbol{\chi}]\!]$, and let $\widetilde{\mathbb{J}} := \mathbb{J}(S)$, where $\mathbb{J}(S)$ is defined as in §4. It follows from Lemma 4.7 that $\mathsf{density}\left(\widetilde{\mathbb{J}}\right) = 1$. For any $m \in \mathbb{N}$, let $\boldsymbol{\chi}^m := \boldsymbol{\chi}\circ\Phi^m$.

LEMMA 7.3. *Let $j \in \widetilde{\mathbb{J}}$, with $j = m + p^r\cdot h$. Then $\boldsymbol{\chi}\circ\Phi^j = \boldsymbol{\chi}^m\circ\Phi^{h'\cdot\langle\!\langle \boldsymbol{\chi}^m \rangle\!\rangle}$, where $h' = p^s\cdot h$ for some $s \geq 0$.*

Proof. Apply Lemma 4.8 to observe that $\Phi^j = \Phi^m\circ\Phi^{h\cdot(p^r)}$. Thus,

$$\boldsymbol{\chi}\circ\Phi^j \quad = \quad \boldsymbol{\chi}\circ\Phi^m\circ\Phi^{h\cdot(p^r)} \quad = \quad \boldsymbol{\chi}^m\circ\Phi^{h\cdot(p^r)}.$$

By definition, r is such that $m < p^{r-1}$ and $[\![\boldsymbol{\chi}]\!] < p^{r-1}$. Thus,

$$[\![\boldsymbol{\chi}^m]\!] \quad = \quad [\![\boldsymbol{\chi}]\!] + m \quad < \quad p^{r-1} + p^{r-1} \quad \leq \quad p^r.$$

Now, let $s := r - \log_p [\![\boldsymbol{\chi}^m]\!]$, and let $h' := p^s\cdot h$. Then $h\cdot(p^r) = h'\cdot\langle\!\langle \boldsymbol{\chi}^m \rangle\!\rangle$, so that $\Phi^{h\cdot(p^r)} = \Phi^{h'\cdot\langle\!\langle \boldsymbol{\chi}^m \rangle\!\rangle}$. □

Proof of Theorem 7.1. We will use Lemma 3.2.

'$\Longleftarrow$' For any $m \in \mathbb{N}$, let $r(m) := \left\lceil \log_p\left(\max\left\{m,\ [\![\boldsymbol{\chi}]\!]\right\}\right)\right\rceil + 1$, and define

$$\widetilde{\mathbb{J}}_m \quad := \quad \left\{m + p^{r(m)}h\ ;\ h \in \mathbb{N}\right\}. \tag{21}$$

It follows that:

$$\widetilde{\mathbb{J}} \quad = \quad \bigcup_{m\in\mathbb{N}} \widetilde{\mathbb{J}}_m. \tag{22}$$

If $j \;=\; m + p^{r(m)}h$ is an element of $\widetilde{\mathbb{J}}_m$, then Lemma 7.3 says $\boldsymbol{\chi}\circ\Phi^j \;=\; \boldsymbol{\chi}^m \circ \Phi^{h'\cdot\langle\!\langle \boldsymbol{\chi}^m\rangle\!\rangle}$, for some $h' \geq h$. Now, μ is Lucas mixing, so find a subset $\widetilde{\mathbb{H}}_m \subset \mathbb{N}$ of density one with $\displaystyle\lim_{\widetilde{\mathbb{H}}_m\ni h\to\infty} \left\langle \boldsymbol{\chi}^m \circ \Phi^{h\cdot\langle\!\langle \boldsymbol{\chi}^m\rangle\!\rangle},\ \mu\right\rangle \;=\; 0$. Define:

$$\begin{aligned}
\mathbb{H}_m &:= \left\{h\in\widetilde{\mathbb{H}}_m\ ;\ \left|\left\langle \boldsymbol{\chi}^m \circ \Phi^{h\cdot\langle\!\langle \boldsymbol{\chi}^m\rangle\!\rangle},\ \mu\right\rangle\right| \;\leq\; \frac{1}{m}\right\}, \\
\mathbb{J}_m &:= \left\{m + p^{r(m)}h\ ;\ h\in\mathbb{H}_m\right\}, \qquad (23)\\
\text{and } \mathbb{J} &:= \bigcup_{m\in\mathbb{N}} \mathbb{J}_m. \qquad (24)
\end{aligned}$$

Claim 1: $\mathsf{density}\,(\mathbb{J}) = 1$.

Proof. For any $m\in\mathbb{N}$, there is some K such that $\mathbb{H}_m = \widetilde{\mathbb{H}}_m \cap [K...\infty)$. Thus, $\mathsf{rel\,density}\left[\mathbb{H}_m/\widetilde{\mathbb{H}}_m\right] = 1$. Thus, $\mathsf{density}\,(\mathbb{H}_m) = \mathsf{density}\left(\widetilde{\mathbb{H}}_m\right) = 1$. Compare (21) and (23) to see that $\mathsf{rel\,density}\left[\mathbb{J}_m/\widetilde{\mathbb{J}}_m\right] = 1$. Then compare (22) and (24) to see that $\mathsf{rel\,density}\left[\mathbb{J}/\widetilde{\mathbb{J}}\right] = 1$. Thus, $\mathsf{density}\,(\mathbb{J}) = \mathsf{density}\left(\widetilde{\mathbb{J}}\right) = 1$. ◇ `Claim 1`

Claim 2: $\displaystyle\lim_{\mathbb{J}\ni j\to\infty} \left\langle \boldsymbol{\chi}\circ\Phi^j,\ \mu\right\rangle \;=\; 0$.

Proof. Fix $\epsilon > 0$. Let M be large enough that $\frac{1}{M} < \epsilon$. For all $m\in\mathbb{N}$ with $m < M$, find H_m such that, if $h\in\widetilde{\mathbb{H}}_m$ and $h > H_m$, then $\left|\left\langle \boldsymbol{\chi}^m \circ \Phi^{h\cdot\langle\!\langle \boldsymbol{\chi}^m\rangle\!\rangle},\ \mu\right\rangle\right| \;<\; \epsilon$. Let $J_m := m + 2^{r(m)}\cdot H_m$. Thus, if $j \;=\; m + 2^{r(m)}\cdot h$ is an element of $\mathbb{J}_m$, and $j \;>\; J_m$, then we must have $h > H_m$, so that $\left|\left\langle \boldsymbol{\chi}\circ\Phi^j,\ \mu\right\rangle\right| \;=\; \left|\left\langle \boldsymbol{\chi}^m \circ \Phi^{h\cdot\langle\!\langle \boldsymbol{\chi}^m\rangle\!\rangle},\ \mu\right\rangle\right| \;<\; \epsilon$.

Now let $J \;:=\; \displaystyle\max_{1\leq m\leq M} J_m$. Thus, for all $j\in\mathbb{J}$, if $j > J$, then either $j\in\mathbb{J}_m$ for some $m\leq M$, in which case $\left|\left\langle \boldsymbol{\chi}\circ\Phi^j,\ \mu\right\rangle\right| \;<\; \epsilon$ by construction of J, or $j\in\mathbb{J}_m$ for some $m > M$, in which case

$$\left|\left\langle \boldsymbol{\chi}\circ\Phi^j,\ \mu\right\rangle\right| \quad \underset{(*)}{<} \quad \frac{1}{m} \quad < \quad \frac{1}{M} \quad \underset{(\dagger)}{<} \quad \epsilon.$$

Here, $(*)$ follows by definition of $\mathbb{H}_m$, and $(\dagger)$ follows by definition of M. ◇ `Claim 2`

Lemma 3.2 and Claims 1 and 2 imply that Φ asymptotically randomizes μ.

'$\Longrightarrow$' Suppose μ was not weakly harmonically mixing. Thus, there is some $\boldsymbol{\chi}\in\widehat{\mathcal{A}^{\mathbb{Z}}}$ and some subset $\mathbb{H}\subset\mathbb{N}$ of density $\delta > 0$ such that $\displaystyle\limsup_{\mathbb{H}\ni h\to\infty} \left|\left\langle \boldsymbol{\chi}\circ\Phi^{h\cdot\langle\!\langle \boldsymbol{\chi}\rangle\!\rangle},\ \mu\right\rangle\right| \;>\; 0$. But $\boldsymbol{\chi}\circ\Phi^{h\cdot\langle\!\langle \boldsymbol{\chi}\rangle\!\rangle} \;=\; \boldsymbol{\chi}\circ\Phi^{p^r\cdot h}$ (where $r = \left\lceil \log_p [\![\boldsymbol{\chi}]\!]\right\rceil$). Hence, if $\mathbb{J} \;:=\; p^r\cdot\mathbb{H}$, then $\mathsf{density}\,(\mathbb{J}) = p^{-r}\cdot\delta \;>\; 0$, and $\displaystyle\limsup_{\mathbb{J}\ni j\to\infty} \left|\left\langle \boldsymbol{\chi}\circ\Phi^j,\ \mu\right\rangle\right| \;=\; \limsup_{\mathbb{H}\ni h\to\infty} \left|\left\langle \boldsymbol{\chi}\circ\Phi^{h\cdot\langle\!\langle \boldsymbol{\chi}\rangle\!\rangle},\ \mu\right\rangle\right| \;>\; 0$. But then Lemma 3.2 implies that Φ cannot randomize μ. □

8. *Randomization of Zero-Entropy Measures*

Of the probability measures which are asymptotically randomized by LCA, every known example has positive entropy. However, we'll show that positive entropy is *not* necessary, by constructing a class of zero-entropy measures which are Lucas mixing, and thus (by Theorem 7.1) randomized by $\Phi = 1 + \sigma$.

For both efficiency and lucidity, we will employ probabilistic language. Let $(\Omega, \mathcal{B}, \rho)$ be an abstract probability space (called the *sample space*). If $(\mathbf{X}, \mathcal{X})$ is any measurable space, then an ($\mathbf{X}$-valued) *random variable* is a measurable function $f : \Omega \longrightarrow \mathbf{X}$. In particular, a *random sequence* is a measurable function $\mathbf{a} : \Omega \longrightarrow \mathcal{A}^{\mathbb{Z}}$. By convention, we suppress the argument of random variables. Thus, if $\mathbf{a}, \mathbf{b}, \mathbf{c}$ are random sequences, then the equation "$\mathbf{a} + \mathbf{b} = \mathbf{c}$" means "$\mathbf{a}(\omega) + \mathbf{b}(\omega) = \mathbf{c}(\omega)$, for ρ-almost all $\omega \in \Omega$."

If $f : \Omega \longrightarrow \mathbf{X}$ is a random variable, and $\mathbf{U} \subset \mathbf{X}$, then "$\mathsf{Prob}\left[f \in \mathbf{U}\right]$" denotes $\rho\left[f^{-1}(\mathbf{U})\right]$. If $g : \Omega \longrightarrow \mathbf{Y}$ is another random variable, then f and g are *independent* if, for any measurable $\mathbf{U} \subset \mathbf{X}$ and $\mathbf{V} \subset \mathbf{Y}$, $\mathsf{Prob}\left[f \in \mathbf{U} \text{ and } g \in \mathbf{V}\right] = \mathsf{Prob}\left[f \in \mathbf{U}\right] \cdot \mathsf{Prob}\left[g \in \mathbf{V}\right]$ —i.e. $\rho\left[f^{-1}(\mathbf{U}) \cap g^{-1}(\mathbf{V})\right] = \rho\left[f^{-1}(\mathbf{U})\right] \cdot \rho\left[g^{-1}(\mathbf{V})\right]$. The *distribution* of f is the probability measure $\mu := f(\rho)$ on $(\mathbf{X}, \mathcal{X})$; we then say that f is a μ-*random variable*. Thus, every random variable determines a probability measure on its range. However, given a measure μ, we can construct infinitely many independent μ-random variables.

Let $\mathcal{A} := \mathbb{Z}_{/2}$ and $\mu \in \mathcal{M}(\mathcal{A}^{\mathbb{Z}})$, and consider a μ-random sequence $\mathbf{a} \in \mathcal{A}^{\mathbb{Z}}$. We say μ has *independent random dyadic increments* (IRDI) if, for any $n \in \mathbb{N}$, and all $m \in [1...2^n]$, $a_{m+2^n} = a_m + d^n_m$, where $d^n_1, \ldots, d^n_{2^n}$ are independent $\mathcal{A}$-valued random variables. If $d^n_1, \ldots, d^n_{2^n}$ have distributions $\delta^n_1, \ldots, \delta^n_{2^n}$, then μ has *lower decay rate* $\alpha \in (0, 1)$ if there is some $L > 0$ such that, for all $n \geq L$, and all $m \in [1...2^n]$, $\alpha^n \leq \delta^n_m\{1\}$.

Proposition 8.1. *If μ has IRDI with lower decay rate $\alpha > \frac{1}{\sqrt{2}}$, then μ is Lucas Mixing.*

Proof. Let $\boldsymbol{\chi} \in \widehat{\mathcal{A}^{\mathbb{Z}}}$ be a nontrivial character. We seek $\mathbb{H} \subset \mathbb{N}$ with $\mathsf{density}\,(\mathbb{H}) = 1$, such that $\lim_{\mathbb{H} \ni h \to \infty} \left\langle \boldsymbol{\chi} \circ \Phi^{h \cdot \langle\!\langle \boldsymbol{\chi} \rangle\!\rangle},\ \mu \right\rangle = 0$.

If $n \in \mathbb{N}$, let $I = I(n) := \lceil \log_2(n) \rceil$, and suppose n has binary expansion $\{n^{(i)}\}_{i=0}^{I}$. Let $\mathbb{I}(n) := \left\{j \in [0...I]\ ;\ n^{(j)} = 1\right\}$. Let $\epsilon > 0$ be small, and define:

$$\mathbb{H} \quad := \quad \left\{h \in \mathbb{N}\ ;\ \#\left(\mathbb{I}(h)\right) \quad \geq \quad \frac{1}{2} I(h) - \epsilon\right\}.$$

Then $\mathsf{density}\,(\mathbb{H}) = 1$. Suppose $n \in \mathbb{H}$ is large; let $\mathbb{I} := \mathbb{I}(n)$ and $I := I(n)$. Assume I is large (in particular, $I > L$).

Now, $\alpha > \frac{1}{\sqrt{2}}$, so find β such that $\frac{1}{\alpha} < \beta < \sqrt{2}$. Define

$$M \quad := \quad \#\left(\mathbb{I}\right) - 1 \quad \geq \quad \frac{1}{2} I - \epsilon - 1 \quad \underset{(*)}{>} \quad \log_2(\beta) I, \tag{25}$$

where $(*)$ is because $\log_2(\beta) < \frac{1}{2}$ and I is large, while ϵ is small.

Suppose $\mathbb{I} = \{i_1 < i_2 < \ldots < i_{M+1} = I\}$. Let $\boldsymbol{\xi}_0 := \boldsymbol{\chi}$, and for each $m \in [0...M]$, define $\boldsymbol{\xi}_{m+1} := \boldsymbol{\xi}_m \otimes \left(\boldsymbol{\xi}_m \circ \sigma^{L_i}\right)$, where $L_i := 2^{i_m} \cdot \langle\!\langle \chi \rangle\!\rangle$. Thus, $\boldsymbol{\chi} \circ \Phi^{n \cdot \langle\!\langle \chi \rangle\!\rangle} = \boldsymbol{\xi}_{M+1}$.

Let $r := \mathsf{rank}\left[\boldsymbol{\chi}\right]$. Then for all $m \in [1...M+1]$, $\mathsf{rank}\left[\boldsymbol{\xi}_m\right] = 2^m \cdot r$. In particular, define

$$R \quad := \quad \mathsf{rank}\left[\boldsymbol{\xi}_M\right] \quad = \quad 2^M \cdot r \quad \underset{(*)}{>} \quad \beta^I \cdot r. \tag{26}$$

where $(*)$ is by equation (25). Thus, $\boldsymbol{\xi}_M = \bigotimes_{x\in\mathbb{X}} \xi_x$, where $\mathbb{X} \subset \mathbb{Z}$ is a subset with $\#(\mathbb{X}) = R$. Thus, if $\mathbf{a} \in \mathcal{A}^{\mathbb{Z}}$ is a μ-random sequence, then

$$\begin{aligned}\boldsymbol{\xi}_{M+1}(\mathbf{a}) &= \boldsymbol{\xi}_M(\mathbf{a}) \cdot \left(\boldsymbol{\xi}_M \circ \sigma^{2^I}(\mathbf{a})\right) = \prod_{x\in\mathbb{X}} \xi_x(a_x)\cdot \xi_x\left(a_{x+2^I}\right) \\ &= \prod_{x\in\mathbb{X}} \xi_x\left(a_x + a_{x+2^I}\right) = \prod_{x\in\mathbb{X}} \xi_x\left(d_x^I\right), \end{aligned} \tag{27}$$

where $\{d_x^I\}_{x\in\mathbb{X}}$ are independent random dyadic increments. If d_x^I has distribution δ_x^I, then

$$\mathbb{E}_{\delta_x^I}\left[\xi_x\left(d_x^I\right)\right] = \delta_x^I\{0\} - \delta_x^I\{1\} = 1 - 2\delta_x^I\{1\} \underset{(*)}{\leq} 1 - 2\cdot\alpha^I = \frac{2\alpha^{-I}-1}{2\alpha^{-I}}. \tag{28}$$

Here, $(*)$ is because μ has lower decay rate α, so $\delta_x^I\{1\} \geq \alpha^I$ (assuming $I \geq L$).

Thus, $\langle \mu,\ \boldsymbol{\chi}\circ\Phi^n\rangle \underset{(\ddagger)}{=} \mathbb{E}_\mu\left[\prod_{x\in\mathbb{X}} \xi_x\left(d_x^I\right)\right] \underset{(*)}{=} \prod_{x\in\mathbb{X}} \mathbb{E}_{\delta_x^I}\left[\xi_x\left(d_x^I\right)\right] \underset{(\dagger)}{\leq} \left(\frac{2\alpha^{-I}-1}{2\alpha^{-I}}\right)^R.$

Here, $(\ddagger)$ is by equation (27), $(*)$ is because $\{d_x^I\}_{x\in\mathbb{X}}$ are independent, and $(\dagger)$ is by equation (28) and because $\#(\mathbb{X}) = R$.

Thus,

$$\begin{aligned}\log\left|\langle\mu,\ \boldsymbol{\chi}\circ\Phi^n\rangle\right| &\leq R\cdot\left[\log\left(2\alpha^{-I}-1\right) - \log(2\alpha^{-I})\right] \underset{(*)}{\leq} -R\cdot\log'\left(2\alpha^{-I}\right) \\ &= \frac{-R}{2\alpha^{-I}} \underset{(\dagger)}{<} \frac{-\beta^I r}{2\alpha^{-I}} = -\frac{r}{2}\,(\alpha\beta)^I.\end{aligned}$$

Here, $(*)$ is because log is a decreasing function, and $(\dagger)$ is by equation (26).

But $\beta > \frac{1}{\alpha}$, so $\alpha\beta > 1$. Thus, $\lim_{\mathbb{H}\ni h\to\infty} \log\left|\left\langle\mu,\ \boldsymbol{\chi}\circ\Phi^{h\cdot\langle\!\langle\boldsymbol{\chi}\rangle\!\rangle}\right\rangle\right| = -\frac{r}{2}\lim_{I\to\infty}(\alpha\beta)^I = -\infty.$
Hence $\lim_{\mathbb{H}\ni h\to\infty}\left|\left\langle\mu,\ \boldsymbol{\chi}\circ\Phi^{h\cdot\langle\!\langle\boldsymbol{\chi}\rangle\!\rangle}\right\rangle\right| = 0.$ □

Suppose $\mu \in \mathcal{M}(\mathcal{A}^{\mathbb{Z}})$ has independent random dyadic increments; for any $n \in \mathbb{N}$, and all $m \in [1...2^n]$, let $\delta_1^n, \ldots, \delta_{2^n}^n$ be the dyadic increment distributions, as before. Then μ has ***upper decay rate*** $\alpha \in (0,1)$ if there are constants $L_1, K > 0$ such that, for all $n \geq L_1$, and all $m \in [1...2^n]$, $\delta_m^n\{1\} \leq K\cdot\alpha^n$.

Proposition 8.2. *If μ has IRDI with upper decay rate $\alpha < 1$, then $h(\mu) = 0$.*

Proof. Let $L_1, K > 0$ be as above. Assume without loss of generality that $K > 4$. Let $L_2 := \dfrac{-\log_2(K)-1}{\log_2(\alpha)}$. Let $L := \max\{L_1, L_2\}$.

For any $n \in \mathbb{N}$, and $m \in [1...2^n]$, let δ_m^n be as above. The entropy of δ_m^n is defined:

$$H(\delta_m^n) := -\delta_m^n\{0\}\log_2(\delta_m^n\{0\}) - \delta_m^n\{1\}\log_2(\delta_m^n\{1\}) \tag{29}$$

Claim 1: *There exists $c_1 > 0$ such that, if $n > L$ and $m \in [1...2^n]$, then $H(\delta_m^n) < c_1 n\cdot\alpha^n$.*

Proof. $\alpha < 1$, so $\log_2(\alpha) < 0$; Thus, if $n \geq L_2$, then $n\log_2(\alpha) \leq L_2 \log_2(\alpha)$. Thus,

$$\begin{aligned}\log_2(K\alpha^n) &= \log_2(K) + n\log_2(\alpha) \quad \leq \quad \log_2(K) + L_2\log_2(\alpha) \\ &= \log_2(K) - \log_2(K) - 1 \quad = \quad -1. \end{aligned} \tag{30}$$

Thus, $\delta_m^n\{1\} \underset{(*)}{\leq} K\alpha^n \underset{(\dagger)}{\leq} \frac{1}{2}$, where $(*)$ is because $n \geq L_1$ and $(\dagger)$ is by equation (30).

But, if $\delta_m^n\{1\} < \frac{1}{2}$ in equation (29), then $H(\delta_m^n)$ decreases as $\delta_m^n\{1\}$ decreases. Hence,

$$\begin{aligned} H(\delta_m^n) &\leq -K\alpha^n \log_2(K\alpha^n) \;-\; (1-K\alpha^n)\log_2(1-K\alpha^n) \\ &< K\alpha^n \underbrace{(nA-k)}_{(*)} \;+\; (1-K\alpha^n)\cdot \underbrace{2K\alpha^n}_{(\dagger)} \quad = \quad K\,(nA+2-k-2K\alpha^n)\cdot\alpha^n \\ &\underset{(\ddagger)}{<} KnA\cdot\alpha^n \quad \underset{(\diamond)}{<} \quad c_1 n\cdot\alpha^n. \end{aligned}$$

Here, $(*)$ is the substitution $k := \log_2(K)$ and $A := -\log_2(\alpha)$; $(\dagger)$ is because, if ϵ is small, then $\log(1-\epsilon) \approx -\epsilon$, thus, $-\log(1-\epsilon) < 2\epsilon$; $(\ddagger)$ is because $2 - k - 2K\alpha^n < 0$ because $k > 2$ because we assume $K > 4$; $(\diamond)$ is where $c_1 := KA > 0$. ◇ `Claim 1`

Let $\mathbf{a} \in \mathcal{A}^{\mathbb{Z}}$ be a μ-random sequence, and fix $n > L$. To compute the conditional entropy $H\left(\mathbf{a}\big|_{(2^n\ldots 2^{n+1}]} \big| \mathbf{a}\big|_{[1\ldots 2^n]}\right)$, recall that, for all $m \in [1...2^n]$, $a_{2^n+m} = a_m + d_m^n$. Thus,

$$\begin{aligned} H\left(\mathbf{a}\big|_{(2^n\ldots 2^{n+1}]} \,\Big|\, \mathbf{a}\big|_{[1\ldots 2^n]}\right) &= H\left(d_1^n, d_2^n, \ldots, d_{2^n}^n\right) \underset{(*)}{=} \sum_{m=1}^{2^n} H(\delta_m^n) \\ &\underset{(\dagger)}{<} 2^n \cdot c_1 n\alpha^n \quad = \quad c_1 n \cdot (2\alpha)^n. \end{aligned} \tag{31}$$

where $(*)$ is because $d_1^n, d_2^n, \ldots, d_{2^n}^n$ are independent random variables with distributions $\delta_1^n, \ldots, \delta_{2^n}^n$, and $(\dagger)$ is by Claim 1. Thus, for any $N > L$,

$$\begin{aligned} H\left(\mathbf{a}\big|_{[1\ldots 2^N]} \,\Big|\, \mathbf{a}\big|_{[1\ldots 2^L]}\right) &= \sum_{n=L}^{N-1} H\left(\mathbf{a}\big|_{(2^n\ldots 2^{n+1}]} \,\Big|\, \mathbf{a}\big|_{[1\ldots 2^n]}\right) \underset{(*)}{<} \sum_{n=L}^{N-1} c_1 n \cdot (2\alpha)^n \\ &< c_1 N \cdot (2\alpha)^L \sum_{n=0}^{N-L-1} (2\alpha)^n \quad = \quad c_1 N \cdot (2\alpha)^L \, \frac{(2\alpha)^{N-L} - 1}{2\alpha - 1} \\ &\leq c_2 N \cdot (2\alpha)^N, \end{aligned} \tag{32}$$

where $(*)$ is by equation (31), and where $c_2 \approx \dfrac{c_1}{2\alpha - 1} > 0$ is another constant.

Thus, if $H_0 := H\left(\mathbf{a}\big|_{[1\ldots 2^L]}\right)$, then

$$H\left(\mathbf{a}\big|_{[1\ldots 2^N]}\right) = H\left(\mathbf{a}\big|_{[1\ldots 2^N]} \,\Big|\, \mathbf{a}\big|_{[1\ldots 2^L]}\right) + H_0 \underset{(*)}{\leq} c_2 N \cdot (2\alpha)^N + H_0, \tag{33}$$

where $(*)$ is by equation (32). Thus,

$$\begin{aligned} h(\mu) &= \lim_{M\to\infty} \frac{1}{M} H\left(\mathbf{a}\big|_{[1\ldots M]}\right) \quad = \quad \lim_{N\to\infty} \frac{1}{2^N} H\left(\mathbf{a}\big|_{[1\ldots 2^N]}\right) \\ &\underset{(*)}{\leq} \lim_{N\to\infty} \frac{c_2 N \cdot (2\alpha)^N + H_0}{2^N} \quad \leq \quad c_2 \lim_{N\to\infty} N\alpha^N \underset{(\dagger)}{=} 0, \end{aligned}$$

where $(*)$ is by equation (33), and $(\dagger)$ is because $|\alpha| < 1$. □

$$\begin{array}{cccccccccccccccccccc}
&&&&&&&&&&&&&&&& r_0^4 & r_1^4 & r_2^4 & r_3^4\dots \\
&&&&&&&& r_0^3 & r_1^3 & r_2^3 & r_3^3 & r_4^3 & r_5^3 & r_6^3 & r_7^3 &&&& \\
&&&& r_0^2 & r_1^2 & r_2^2 & r_3^2 &&&&& r_0^2 & r_1^2 & r_2^2 & r_3^2 &&&& \\
&& r_0^1 & r_1^1 &&& r_0^1 & r_1^1 &&& r_0^1 & r_1^1 &&& r_0^1 & r_1^1 &&& r_0^1 & r_1^1\dots \\
& r_0^0 && r_0^0 && r_0^0 && r_0^0 && r_0^0 && r_0^0 && r_0^0 && r_0^0 && r_0^0 && r_0^0\dots \\
\hline
\dots a_0^\infty & a_1^\infty & a_2^\infty & a_3^\infty & a_4^\infty & a_5^\infty & a_6^\infty & a_7^\infty & a_8^\infty & a_9^\infty & a_{10}^\infty & a_{11}^\infty & a_{12}^\infty & a_{13}^\infty & a_{14}^\infty & a_{15}^\infty & a_{16}^\infty & a_{17}^\infty & a_{18}^\infty & a_{19}^\infty\dots \\
&&&&& \dots & a_0 & a_1 & a_2 & a_3 & a_4 & a_5 & a_6 & a_7 & \dots &&&&&
\end{array}$$

FIGURE 3. The construction of random sequence $\mathbf{a}^\infty$; the approximation of $\mathbf{a}$ as a random translate of $\mathbf{a}^\infty$.

It remains to actually construct a measure with IRDI. Let $0 < \alpha < 1$. For any $n \in \mathbb{N}$, let ρ_n be the probability distribution on $\mathcal{A} = \mathbb{Z}_{/2}$ such that

$$\rho_n\{1\} = \alpha^n \qquad \text{and} \qquad \rho_n\{0\} = 1 - \alpha^n. \tag{34}$$

For each $n \in \mathbb{N}$, we will construct a random sequence $\mathbf{a}^n \in \mathcal{A}^{\mathbb{Z}}$ as follows. First, define $\mathbf{a}^0 := [\dots 0000 \dots]$. Now, suppose, inductively, that we have $\mathbf{a}^n$. Let $r_0^n, r_1^n, \dots, r_{2^n-1}^n$ be a set of 2^n independent $\mathcal{A}$-valued, ρ_n-random variables. Let $\mathbf{r}^n \in \mathcal{A}^{\mathbb{Z}}$ be the random, 2^{n+1}-periodic sequence

$$\mathbf{r}^n \;:=\; [\dots, \underbrace{\overset{\substack{\text{zeroth coordinate}\\ \downarrow}}{0}, 0, \dots, 0}_{2^n}, r_0^n, r_1^n, \dots, r_{2^n-1}^n, \underbrace{0, 0, \dots, 0}_{2^n}, r_0^n, r_1^n, \dots, r_{2^n-1}^n, \dots],$$

and inductively define $\mathbf{a}^{n+1} := \mathbf{a}^n + \mathbf{r}^n$.

Let $\mu_n \in \mathcal{M}(\mathcal{A}^{\mathbb{Z}})$ be the distribution of $\mathbf{a}^n$, and let $\widetilde{\mu}_n := \dfrac{1}{2^n} \displaystyle\sum_{i=1}^{2^n} \sigma^i(\mu_n)$ be the stationary average of μ_n. Finally, let $\mu := \text{wk}^*\text{–}\lim\limits_{n\to\infty} \widetilde{\mu}_n$.

Let μ_∞ be the probability distribution of the random sequence $\mathbf{a}^\infty := \displaystyle\sum_{n=1}^{\infty} \mathbf{r}^n$ (see Fig.3). Then $\mu_\infty = \text{wk}^*\text{–}\lim\limits_{n\to\infty} \mu_n$, and loosely speaking, μ is the 'σ-ergodic average' of μ_∞. Thus, if $\mathbf{a}$ is a μ-random sequence, we can think of $\mathbf{a}$ as obtained by shifting $\mathbf{a}^\infty$ by a random amount. The next lemma describes the structure of $\mathbf{a}^\infty$:

LEMMA 8.3. *Let $M \in \mathbb{N}$ have binary expansion $M = \displaystyle\sum_{n=0}^{\infty} m_n 2^n$. For all $n \geq 0$, let $M_n := \displaystyle\sum_{i=0}^{n-1} m_i 2^i$. Then $a_M^\infty \;=\; \displaystyle\sum_{n=0}^{\infty} m_n \cdot r_{M_n}^n$.* □

For example, suppose $M := 13 = 1 + 4 + 8$; then $m_0 = m_2 = m_3 = 1$ and $m_1 = 0$. Hence, $M_0 = 0,\ M_1 = M_2 = 1$, and $M_3 = 5$. Thus, $a_{13}^\infty \;=\; r_0^0 + r_1^2 + r_5^3$ (see Figure 3).

Think of $\mathbf{a}^\infty$ as being generated by a process of 'duplication with error'. Let $\mathbf{w}^0 := [0]$ be a word of length 1. Suppose, inductively, that we have $\mathbf{w}^n = [w_1 w_2 \dots w_{2^n-1}]$. Let $\widetilde{\mathbf{w}}^n := [\widetilde{w}_1 \widetilde{w}_2 \dots \widetilde{w}_{2^n-1}]$ be an 'imperfect copy' of $\mathbf{w}^n$: for each $m \in [0...2^n)$, $\widetilde{w}_m := w_m + r_m^n$, where $r_0^n, r_1^n \dots, r_{2^n-1}^n$ are the independent ρ_n-distributed variables from before, which act as 'copying errors'. Let $\mathbf{w}^{n+1} := \mathbf{w}^n \widetilde{\mathbf{w}}^n$. Then $\mathbf{a}^\infty$ is the limit of $\mathbf{w}^n$ as $n \to \infty$.

PROPOSITION 8.4. *μ has IRDI, with upper and lower decay rate α.*

Proof. Let $\mathbf{a} \in \mathcal{A}^{\mathbb{Z}}$ be a μ-random sequence, and fix $N \in \mathbb{N}$. By construction, there is some $k \in \mathbb{Z}$ such that $\mathbf{a}$ looks like $\sigma^k(\mathbf{a}^\infty)$ in a neighbourhood around 0. To be precise,

$$\text{For all } m \in \big[0...2^{N+1}\big), \qquad a_m \quad = \quad a^\infty_{k+m}. \tag{35}$$

For example, in Figure 3, let $N = 2$, so that $2^N = 4$; suppose $k = 6$. Thus, $[a_0, a_1, \ldots, a_7] = [a^\infty_6, a^\infty_7, \ldots, a^\infty_{13}]$. Thus, $d^2_0 = a_4 - a_0 = a^\infty_{10} - a^\infty_6 = r^3_2 - r^2_2 = r^3_2 + r^2_2$. More generally:

CLAIM 1: *Let $m \in \big[0...2^N\big)$.*

(a) *There is a set $S(m) := \{(n_0, m_0),\ (n_1, m_1), \ldots, (n_J, m_J)\}$ (for some $J \geq 0$), where $N = n_0 \leq n_1 \leq \cdots \leq n_J$, and where $m_j \in [0...2^{n_j})$ for $\forall\ j \in [0...J]$, such that $d^N_m = r^{n_0}_{m_0} + r^{n_1}_{m_1} + \ldots + r^{n_J}_{m_J}$.*

(b) *If $m' \in \big[0...2^N\big)$, and $m' \neq m$, then $S(m') \cap S(m) = \emptyset$.*

Proof. Let $M := k + m$ and let $\widetilde{M} := k + m + 2^N$. If $M = \sum_{n=0}^{\infty} m_n 2^n$ and $\widetilde{M} = \sum_{n=0}^{\infty} \widetilde{m}_n 2^n$, then Lemma 8.3 says that

$$a^\infty_M \quad = \quad \sum_{n=0}^{\infty} m_n \cdot r^n_{M_n}; \qquad \text{and} \qquad a^\infty_{\widetilde{M}} \;=\; \sum_{n=0}^{\infty} \widetilde{m}_n \cdot r^n_{\widetilde{M}_n}. \tag{36}$$

Let $N_1 \geq N$ be the smallest element of $[N...\infty)$ such that $m_{N_1} = 0$. Hence, $m_n = 1$ for all $n \in [N...N_1)$, and $m_{N_1} = 0$. Note that $\widetilde{M} = M + 2^N$, so binary expansions of M and $\widetilde{M}$ are related as follows:

(A) $m_n = \widetilde{m}_n$ for all $n \in [0...N)$.

(B) Thus, $\widetilde{M}_n = M_n$ for all $n \in [0...N]$.

(C) If $m_N = 0$ then $\widetilde{m}_N = 1$. If $m_N = 1$ then $\widetilde{m}_N = 0$.

(D) $\widetilde{m}_n = 0$ for all $n \in [N...N_1)$ (possibly an empty set), and $\widetilde{m}_{N_1} = 1$.

(E) $m_n = \widetilde{m}_n$ for all $n > N_1$.

Thus,

$$\begin{aligned}
d^N_m \quad &= \quad a_{m+2^N} - a_m \;\underset{(*)}{=}\; a^\infty_{k+m+2^N} - a^\infty_{k+m} \quad = \quad a^\infty_{\widetilde{M}} + a^\infty_M \qquad \pmod 2 \\
&\underset{(\dagger)}{=} \quad \sum_{n=0}^{\infty} \Big(\widetilde{m}_n \cdot r^n_{\widetilde{M}_n} + m_n \cdot r^n_{M_n} \Big) \;\underset{(\mathrm{ab})}{=}\; \sum_{n=N}^{\infty} \Big(\widetilde{m}_n \cdot r^n_{\widetilde{M}_n} + m_n \cdot r^n_{M_n} \Big) \\
&= \quad \underbrace{r^N_{M_N}}_{(\mathrm{bc})} + \sum_{n=N+1}^{N_1-1} m_n \underbrace{r^n_{M_n}}_{(\mathrm{d})} + \underbrace{r^{N_1}_{\widetilde{M}_{N_1}}}_{(\mathrm{d})} + \sum_{n=N_1+1}^{\infty} \underbrace{m_n}_{(\mathrm{e})} \cdot \Big(r^n_{\widetilde{M}_n} + r^n_{M_n} \Big)
\end{aligned} \tag{37}$$

Here, $(*)$ is by equation (35); $(\dagger)$ is by equation (36); (ab) is by (A) and (B); (bc) is by (B) and (C); (d) is by (D), and (e) is by (E).

Now, to see **(a)**, let

$$S(m) \quad := \quad \Big\{(n,m)\ ;\ r^n_m \text{ appears with nonzero coefficient in expression (37)}\Big\}.$$

In particular, $r^N_{M_N}$ appears in (37), so $(n_0, m_0) := (N, M_N)$; thus, $n_0 = N$.

To see **(b)**, suppose $m < m'$; hence $m' = m + i$ for some $i \in \big[1...2^N\big)$.

Let $M' := M + i$ and $\widetilde{M}' := \widetilde{M} + i$. Suppose $M' = \sum_{n=0}^{\infty} m'_n 2^n$ and $\widetilde{M}' = \sum_{n=0}^{\infty} \widetilde{m}'_n 2^n$. Define M'_n, $\widetilde{M}'_n$, and N'_1 analogously. Then, an argument identical to equation (37) yields:

$$d^N_{m'} \quad = \quad r^N_{M'_N} \quad + \sum_{n=N+1}^{N'_1-1} m'_n r^n_{M'_n} \quad + \quad r^{N'_1}_{\widetilde{M}'_{N'_1}} \quad + \sum_{n=N'_1+1}^{\infty} m'_n \cdot \left(r^n_{\widetilde{M}'_n} + r^n_{M'_n}\right) \qquad (38)$$

Now, for all $n \in [N...\infty)$, $\quad M'_n = M_n + i$ and $\widetilde{M}'_n = \widetilde{M}_n + i$ (because $i < 2^N$); thus, $r^n_{M'_n} = r^n_{M_n+i} \notin \{r^n_{M_n}, r^n_{\widetilde{M}_n}\}$ and $r^n_{\widetilde{M}'_n} = r^n_{\widetilde{M}_n+i} \notin \{r^n_{M_n}, r^n_{\widetilde{M}_n}\}$. Thus, every summand of equation (38) is distinct from every summand of equation (37), so $S(m') \cap S(m) = \emptyset$.
◇ `Claim 1`

To see that the random variables $d^N_0, \ldots, d^N_{2^N-1}$ are jointly independent, use Claim 1(a):

$$d^N_0 \ = \sum_{(n,m)\in S(0)} r^n_m, \qquad d^N_1 \ = \sum_{(n,m)\in S(1)} r^n_m, \qquad \ldots\ldots \quad d^N_{2^N-1} \ = \sum_{(n,m)\in S(2^N-1)} r^n_m$$

The random variables $\big\{r^n_m\ ;\ n \in \mathbb{N},\ m \in \big[1...2^N\big]\big\}$ are independent, and Claim 1(b) says $S(0), S(1)\ldots, S(2^N-1)$ are pairwise disjoint; thus $d^N_0, \ldots, d^N_{2^N-1}$ are jointly independent.

Lower Decay Rate: $|\alpha| < 1$, so if N is sufficiently large (e.g. $N > L := -1/\log_2(\alpha)$), then $\alpha^N < \frac{1}{2}$. Suppose $d^N_m \ = \ r^{n_0}_{m_0} + r^{n_1}_{m_1} + \ldots + r^{n_J}_{m_J}$, as in Claim 1(a). For all $j \in [0...J]$, let $P_j := \mathsf{Prob}\left(\sum_{i=j}^{J} r^{n_i}_{m_i} \ = \ 1\right)$. Thus,

$$\begin{aligned}\delta^N_m\{1\} \quad &= \quad P_0 \underset{(\dagger)}{=\!=} \rho_N\{0\}\cdot P_1 \ + \ \rho_N\{1\}\cdot(1-P_1) \quad = \quad (1-\alpha^N)\cdot P_1 \ + \ \alpha^N\cdot(1-P_1) \\ &= \quad \alpha^N \ + \ (1-2\alpha^N)\cdot P_1 \quad \underset{(*)}{\geq} \quad \alpha^N\end{aligned}$$

($\dagger$) is because Claim 1(a) says $n_0 = N$. ($*$) is because $1 - 2\alpha^N > 0$, because $\alpha^N \ < \ \frac{1}{2}$.

Upper Decay Rate: Let $K := \frac{1}{1-\alpha}$. We claim that, for any N and m, $\ \delta^N_m\{1\} \ \leq \ K\alpha^N$.

As before, let $P_j := \mathsf{Prob}\left(\sum_{i=j}^{J} r^{n_i}_{m_i} \ = \ 1\right)$. For any $j \in [1...J)$, we have

$$P_j \ = \ (1-\alpha^{n_j})\cdot P_{j+1} + \alpha^{n_j}\cdot(1-P_{j+1}) \ = \ P_{j+1} + (1-2P_{j+1})\alpha^{n_j} \ \leq \ P_{j+1} + \alpha^{n_j}, \qquad (39)$$

and $P_J = \alpha^{n_J}$. Hence,

$$\delta^N_m\{1\} \quad = \quad P_0 \underset{(*)}{\leq} \alpha^{n_0} + \alpha^{n_1} + \ldots + \alpha^{n_J} \quad \leq \quad \sum_{i=n_0}^{\infty} \alpha^i \quad = \quad \frac{\alpha^{n_0}}{1-\alpha} \quad = \quad K\alpha^{n_0} \underset{(\dagger)}{=\!=} K\alpha^N.$$

Here, ($*$) is obtained by applying equation (39) inductively, and ($\dagger$) is because $n_0 = N$.
□

Thus, if $\frac{1}{\sqrt{2}} < \alpha < 1$, then μ satisfies the conditions of Propositions 8.1 and 8.2, so μ is a zero-entropy, Lucas mixing measure. Hence, $1 + \sigma$ asymptotically randomizes μ.

REFERENCES

[Bré99] P. Brémaud. *Markov Chains: Gibbs fields, Monte Carlo Simulation, and Queues.* Springer, 1999.

[FMMN00] Pablo A. Ferrari, Alejandro Maass, Servet Martínez, and Peter Ney. Cesàro mean distribution of group automata starting from measures with summable decay. *Ergodic Theory Dynam. Systems*, 20(6):1657–1670, 2000.

[Kit98] Bruce Kitchens. *Symbolic dynamics: one-sided, two-sided, and countable state Markov shifts.* Springer, New York, 1998.

[KS80] Ross Kindermann and J. Laurie Snell. *Markov Random Fields and their Applications.* American Mathematical Society, Providence, Rhode Island, 1980.

[Lin84] Doug Lind. Applications of ergodic theory and sofic systems to cellular automata. *Physica D*, 10:36–44, 1984.

[LM95] Doug Lind and Brian Marcus. *An Introduction to Symbolic Dynamics and Coding.* Cambridge UP, New York, 1995.

[MHM03] Alejandro Maass, Bernard Host, and Servet Martínez. Uniform Bernoulli measure in dynamics of permutative cellular automata with algebraic local rules. *Discrete & Continuous Dyn. Sys.*, 9(6):1423–1446, November 2003.

[MM98] Alejandro Maass and Servet Martínez. On Cesàro limit distribution of a class of permutative cellular automata. *Journal of Statistical Physics*, 90(1-2):435–452, 1998.

[MM99] Alejandro Maass and Servet Martínez. Time averages for some classes of expansive one-dimensional cellular automata. In Eric Goles and Servet Martinez, editors, *Cellular Automata & Complex Systems*, pages 37–54. Kluwer Academic Publishers, Dordrecht, 1999.

[MMPY06] Alejandro Maass, Servet Martínez, Marcus Pivato, and Reem Yassawi. Asymptotic randomization of subgroup shifts by linear cellular automata. to appear in *Ergodic Theory & Dynamical Systems*, 26, 2006.

[Pet89] Karl Petersen. *Ergodic Theory.* Cambridge University Press, New York, 1989.

[PY02] Marcus Pivato and Reem Yassawi. Limit measures for affine cellular automata. *Ergodic Theory Dynam. Systems*, 22(4):1269–1287, 2002.

[PY04] Marcus Pivato and Reem Yassawi. Limit measures for affine cellular automata. II. *Ergodic Theory Dynam. Systems*, 24(6):1961–1980, 2004.